

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
5 July 2001 (05.07.2001)

PCT

(10) International Publication Number
WO 01/48183 A3

(51) International Patent Classification⁷: **C12N 15/10**,
9/22, C07K 14/435, C12N 15/66, 15/70, 1/00

(74) Agent: **BAYLISS, Geoffrey, Cyril**; Boulton Wade Tennant,
Verulam Gardens, 70 Gray's Inn Road, London WC1X
8BT (GB).

(21) International Application Number: **PCT/EP00/13149**

(22) International Filing Date:
22 December 2000 (22.12.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
9930691.2 24 December 1999 (24.12.1999) GB

(71) Applicant (for all designated States except US): **DEV-
GEN NV** [BE/BE]; Technologiepark 9, B-9052 Zwij-
naarde (BE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **PLAETINCK,**
Geert [BE/BE]; Pontstraat 16, B-9820 Merelbeke (BE).
MORTIER, Katherine [BE/BE]; Paddenhoek 20, B-9830
St.-Martens Latem (BE). **LISSENS, Ann** [BE/BE]; Tiens-
esteenweg 137, B-3010 Kessel-Lo (BE). **BOGAERT,**
Thierry [BE/BE]; Wolvendreef 26g, B-8500 Kortrijk
(BE).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

(88) Date of publication of the international search report:
6 December 2001

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: IMPROVEMENTS RELATING TO DOUBLE-STRANDED RNA INHIBITION

(57) Abstract: There are described ways of improving the efficiency of double stranded RNA inhibition as a method of inhibiting gene expression in nematode worms such as *C. elegans*. In particular, the invention relates to the finding that changes in the genetic background of *C. elegans* result in increased sensitivity to double-stranded RNA inhibition.

BEST AVAILABLE COPY



WO 01/48183 A3

IMPROVEMENTS RELATING TO DOUBLE-STRANDED RNA
INHIBITION

5 The present invention is concerned with ways of
improving the efficiency of double stranded RNA
inhibition as a method of inhibiting gene expression
in nematode worms such as *C. elegans*. In particular,
the invention relates to the finding that the
susceptibility of nematode worms such as *C. elegans* to
10 double stranded RNA inhibition is affected by changes
in the genetic background of the worms.

It has recently been described in Nature Vol 391,
pp.806-811, February 98, that introducing double
stranded RNA into a cell results in potent and
15 specific interference with expression of endogenous
genes in the cell, which interference is substantially
more effective than providing either RNA strand
individually as proposed in antisense technology. This
specific reduction of the activity
20 of the gene was also found to occur in the nematode
worm *Caenorhabditis elegans* (*C. elegans*) when the RNA
was introduced into the genome or body cavity of the
worm.

The present inventors have utilized the double
25 stranded RNA inhibition technique and applied it
further to devise novel and inventive methods of (i)
assigning functions to genes or DNA fragments which
have been sequenced in various projects, such as, for
example, the human genome project and which have yet
30 to be accorded a particular function, and (ii)
identifying DNA responsible for conferring a
particular phenotype. Such methods are described in
the applicant's co-pending application number WO
00/01846. Processes for introducing RNA into a living
35 cell, either *in vivo* or *ex vivo*, in order to inhibit
expression of a target gene in that cell are

additionally described in WO 99/32619.

Several different experimental approaches can be used to introduce double-stranded RNA into nematode worms in order to achieve RNA interference *in vivo*.

5 One of the most straightforward approaches is simple injection of double-stranded RNA into a body cavity. A more elegant solution is to feed the nematodes on food organisms, generally bacteria, which express a double stranded RNA of the appropriate sequence,
10 corresponding to a region of the target gene.

The present inventors have now determined that the phenomenon of RNA interference in nematodes following ingestion of food organisms capable of expressing double-stranded RNA is dependent both on
15 the nature of the food organism and on the genetic background of the nematodes themselves. These findings may be exploited to provided improved methods of double-stranded RNA inhibition.

Therefore, according to a first aspect of the
20 present invention there is provided a method of inhibiting expression of a target gene in a nematode worm comprising feeding to said nematode worm a food organism which is capable of producing a double-stranded RNA structure having a nucleotide sequence
25 substantially identical to a portion of said target gene following ingestion of the food organism by the nematode, wherein the nematode has a non wild-type genetic background selected to provide increased sensitivity to RNA interference as compared to wild
30 type.

Caenorhabditis elegans is the preferred nematode worm for use in the method of the invention although the method could be carried out with other nematodes and in particular with other microscopic nematodes,
35 preferably microscopic nematodes belonging to the genus *Caenorhabditis*. As used herein the term "microscopic" nematode encompasses nematodes of

approximately the same size as *C. elegans*, being of the order 1mm long in the adult stage. Microscopic nematodes of this approximate size can easily be grown in the wells of a multi-well plate of the type
5 generally used in the art to perform mid- to high-throughput screening.

It is an essential feature of this aspect of the invention that the nematode has a non wild-type genetic background which confers greater sensitivity
10 to RNA interference phenomena (abbreviated herein to RNAi) as compared to the equivalent wild type nematodes. As illustrated in the accompanying examples, introduction of double-stranded RNA (abbreviated herein to dsRNA) into a non wild-type
15 strain according to the invention results in greater inhibition of expression of the target gene. Depending on the nature of the target gene, this greater level of inhibition may be detectable at the phenotypic level as a more pronounced phenotype.

The nematode having non wild-type genetic background may, advantageously, be a mutant strain. Mutations which have the effect of increasing susceptibility of the nematode to RNAi may, for
20 example, affect the stability of dsRNA or the kinetics of dsRNA turnover within cells of the worm or the rate of uptake of dsRNA synthesised by a food organism. Suitable mutant strains include mutant strains
25 exhibiting knock-out or loss-of-function mutations in one or more genes encoding proteins involved in RNA synthesis, RNA degradation or the regulation of these
30 processes.

In one preferred embodiment, the nematode is a mutant strain, more preferably a mutant *C. elegans*, which exhibits reduced activity of one or more
35 nucleases compared to wild-type. Suitable strains include mutant strains exhibiting knock-out or loss-of-function mutations in one or more genes encoding

nucleases, such as RNases. A particularly preferred example is the *nuc-1* strain. This mutant *C. elegans* strain is known per se in the art.

5 In a second preferred embodiment, the nematode is a mutant strain, more preferably a mutant *C. elegans*, which exhibits increased gut uptake compared to wild-type. Particularly preferred examples of such strains are the so-called *C. elegans* gun mutants described herein. In a still further embodiment, the nematode
10 may be a transgenic worm comprising one or more transgenes which increase gut uptake relative to wild-type.

The term "increased gut uptake" as used herein is taken to mean increased uptake of foreign particles
15 from the gut lumen and may encompass both increased gut permeability and increased gut molecular transport compared to wild-type *C. elegans*.

C. elegans feeds by taking in liquid containing its food (e.g. bacteria). It then spits out the
20 liquid, crushes the food particles and internalises them into the gut lumen. This process is performed by the muscles of the pharynx. The process of taking up liquid and subsequently spitting it out is called pharyngeal pumping. Once the food particles have been
25 internalised via pharyngeal pumping their contents must cross the gut itself in order to reach target sites in the worm. There are multiple factors which effect the uptake of compounds from the gut lumen to the surrounding tissues. These include the action of
30 multi-drug resistance proteins, multi-drug resistance related proteins and the P450 cytochromes as well as other enzymes and mechanisms available for transport of molecules through the gut wall.

C. elegans mutants which exhibit increased uptake
35 of foreign molecules through the gut may be obtained from the *C. elegans* mutant collection at the C.

C. elegans Genetic Center, University of Minnesota, St Paul, Minnesota, or may be generated by standard methods. Such methods are described by Anderson in Methods in Cell Biology, Vol 48, "C. elegans: Modern biological analysis of an organism" Pages 31 to 58. Several selection rounds of the PCR technique can be performed to select a mutant worm with a deletion in a desired gene. Alternatively, a population of worms could be subjected to random mutagenesis and worms exhibiting the desired characteristic of increased gut uptake selected using a phenotypic screen, such as the dye uptake method described herein.

As an alternative to mutation, transgenic worms may be generated with the appropriate characteristics. Methods of preparing transgenic worms are well known in the art and are particularly described by Craig Mello and Andrew Fire, Methods in Cell Biology, Vol 48, Ed. H.F. Epstein and D.C. Shakes, Academic Press, pages 452-480.

Worms exhibiting the desired characteristics of increased gut uptake can be identified using a test devised by the inventors based on uptake of a marker precursor molecule which is cleaved by the action of enzymes present in the gut lumen to generate a marker molecule which produces a detectable signal, such as fluorescence. A suitable marker precursor molecule is the fluorescent dye precursor BCECF-AM available from Molecular Probes (Europe BV), Netherlands. This dye only becomes fluorescent when cleaved by esterases and maintained at a pH above 6. The pH of the gut lumen is usually 5 or below. Thus, any BCECF-AM taken up through the pharynx into the gut lumen is not fluorescent until cleaved and the cleaved portion has entered the cells surrounding the lumen which are at a higher pH. Thus, this dye is able to quickly identify mutant or otherwise modified worms which have increased gut transport or permeability. There is a

gradual increase in fluorescence in the tissues surrounding the gut while the gut lumen remains dark. The fluorescence can be detected at an excitation wavelength of 485 nm and an emission wavelength of 530 nm.

Specific examples of gun mutant strains isolated using this procedure which may be used in the method of the invention are strains bg77, bg84, bg85 and bg86, although it is to be understood that the invention is in no way limited to the use of these specific strains. The *C. elegans* gun mutant strain bg85 was deposited on 23 December 1999 at the BCCM/LMG culture collection, Laboratorium Voor Microbiologie, Universiteit Gent, K. L. Ledeganckstraat 35, B-9000, Gent, Belgium under accession number LMBP 5334CB. The phrase "the bg85 mutation" as used herein refers to the specific mutation(s) present in the bg85 strain which is/are responsible for conferring the gun phenotype.

It is also within the scope of the invention to use a non wild-type nematode strain, preferable a *C. elegans* strain, having multiple mutations which affect sensitivity to RNAi. A preferred type of multiple mutant is one having at least one mutation which results in reduced nuclease activity compared to wild type and at least one mutation which results in increased gut uptake compared to wild type. An example of such a mutant is a *C. elegans* strain having the *nuc-1* mutation and at least one further gun mutation. As exemplified herein, double mutants having the *nuc-1* mutation and a gun mutation exhibit enhanced sensitivity to RNAi as compared to either *nuc-1* or gun single mutants.

For the avoidance of doubt, where particular characteristics or properties of nematode worms are described herein by relative terms such as "enhanced"

or "increased" or "decreased" this should be taken to mean enhanced, increased or decreased relative to wild-type nematodes. In the case of *C. elegans*, wild-type is defined as the N2 Bristol strain which is well known to workers in the *C. elegans* field and has been extremely well characterised (see Methods in Cell Biology, Volume 48, *Caenorhabditis elegans*: Modern biological analysis of an organism, ed. by Henry F. Epstein and Diane C. Shakes, 1995 Academic Press; The nematode *Caenorhabditis elegans*, ed. by William Wood and the community of *C. elegans* researchers., 1988, Cold Spring Harbor Laboratory Press; *C. elegans* II, ed. by Donald L. Riddle, Thomas Blumenthal, Barbara J. Meyer and James R. Priess, 1997, Cold Spring Harbor Laboratory Press). The N2 strain can be obtained from the *C. elegans* Genetic Center, University of Minnesota, St Paul, Minnesota, USA.

The food organism for use in the above aspect of the invention is preferably a bacterium such as, for example, a strain of *E.coli*. It will, however, be appreciated that any other type of food organism on which nematodes feed and which is capable of producing dsRNA could be used. The food organism may be genetically modified to express a double-stranded RNA of the appropriate sequence, as will be understood with reference to the examples included herein. One convenient way in which this may be achieved in a bacterial food organism is by transforming the bacterium with a vector comprising a promoter or promoters positioned to drive transcription of a DNA sequence to RNA capable of forming a double-stranded structure. Examples of such vectors will be further described below.

The actual step of feeding the food organism to the nematode may be carried out according to procedures known in the art, see WO 00/01846.

Typically the feeding of the food organisms to the nematodes is performed on standard agar plates commonly used for culturing *C. elegans* in the laboratory. However, the step of feeding the food
5 organism to the nematodes may also be carried out in liquid culture, for example in the wells of 96-well microtitre assay plates.

The inventors have further observed that variations in the food organism can result in enhanced
10 *in vivo* RNAi when the food organism is ingested by a nematode worm.

Accordingly, in a further aspect the invention provides a method of inhibiting expression of a target gene in a nematode worm comprising feeding to said
15 nematode worm a food organism capable of producing a double-stranded RNA structure having a nucleotide sequence substantially identical to a portion of said target gene following ingestion of the food organism by the nematode, wherein the food organism carries a
20 modification selected to provide increased expression or persistence of the double-stranded RNA compared to a food organism which does not carry the modification.

The modification present in the food organism can be any modification which results in increased
25 expression of the dsRNA or in increased persistence of the dsRNA. Suitable modifications might include mutations within the bacterial chromosome which affect RNA stability and/or degradation or mutations which have a direct effect on the rate of transcription. In
30 a preferred embodiment, the food organism is an RNase III minus *E. coli* strain, or any other RNase negative strain.

According to a still further aspect of the invention there is provided a method of inhibiting
35 expression of a target gene in a nematode worm comprising introduction of a DNA capable of producing a double-stranded RNA structure having a nucleotide

sequence substantially identical to a portion of said target gene in said nematode, wherein the nematode is one which exhibits increased gut uptake compared to wild type.

5 In addition to exhibiting increased sensitivity to RNAi following feeding with food organisms capable of expressing a dsRNA, nematodes which exhibit increase gut uptake as described herein also show increased uptake of DNA molecules capable of producing
10 double-stranded RNA structures following ingestion into a nematode.

In a preferred embodiment, the DNA is in the form of a vector comprising a promoter or promoters orientated to relative to a sequence of DNA such that
15 they are capable of driving transcription of the said DNA to make RNA capable of forming a double-stranded structure upon binding of an appropriate RNA polymerase to the promoter or promoters.

Several different arrangements of promoters may
20 be used in such a vector. In a first arrangement a DNA fragment corresponding to a region of the target gene is flanked by two opposable polymerase-specific promoters which are preferably identical. Transcription from the opposable promoters produces
25 two complementary RNA strands which can anneal to form an RNA duplex. The plasmid pGN1 described herein is an example of a vector comprising two opposable T7 promoters flanking a multiple cloning site for insertion of a DNA fragment of the appropriate
30 sequence, corresponding to a region of a target gene. pGN8 is an example of a vector derived from pGN1 containing a fragment of the *C. elegans unc-22* gene. In an alternative arrangement, DNA fragments corresponding to a region of the target gene may be
35 placed both in the sense and the antisense orientation downstream of a single promoter. In this case, the sense/antisense fragments are co-transcribed to

generate a single RNA strand which is self-complementary and can therefore form an RNA duplex.

In both of the above arrangements, the polymerase-specific T3, T7 and SP6 promoters, all of which are well known in the art, are useful for driving transcription of the RNA. Expression from these promoters is dependent on expression of the cognate polymerase. Advantageously, the nematode itself may be adapted to express the appropriate polymerase. Expression of the polymerase may be general and constitutive, but could also be regulated under a tissue-specific promoter, an inducible promoter, a temporally regulated promoter or a promoter having a combination of such characteristics. Transgenic *C. elegans* strains harboring a transgene encoding the desired polymerase under the control of an appropriately-regulated promoter can be constructed according to methods known *per se* in the art and described, for example, by Craig Mello and Andrew Fire in *Methods in Cell Biology*, Vol 48, Ed. H. F. Epstein and D. C. Shakes, Academic Press, pp 452-480.

The advantage of adapting the nematode to express the required polymerase is that it is possible to control inhibition of expression of the target gene in a tissue-specific and/or temporally specific manner by placing expression of the polymerase under the control of an appropriately regulated promoter.

Introduction of DNA into nematodes in accordance with the method of the invention can be achieved using a variety of techniques, for example by direct injection into a body cavity or by soaking the worms in a solution containing the DNA. If the DNA is in the form of a vector as described herein, e.g. a plasmid harboring a cloned DNA fragment between two flanking T7 promoters, then dsRNA corresponding to this DNA fragment will be formed in the nematode resulting in down regulation of the corresponding gene. The

introduced DNA can form an extrachromosomal array, which array might result in a more catalytic knock-out or reduction of function phenotype. The DNA might also become integrated into the genome of the nematode, resulting in the same catalytic knock out or reduction of function phenotype, but which is stably transmittable.

In each aspect of the invention, the double-stranded RNA structure may be formed by two separate complementary RNA strands or a single self-complementary strand, as described above. Inhibition of target gene expression is sequence-specific in that only nucleotide sequences corresponding to the duplex region of the dsRNA structure are targeted for inhibition.

It is preferred to use dsRNA comprising a nucleotide sequence identical to a portion of the target gene, although RNA sequences with minor variations such as insertions, deletions and single base substitutions may also be used and are effective for inhibition. It will be readily apparent that 100% sequence identity between the dsRNA and a portion of the target gene is not absolutely required for inhibition and the phrase "substantially identical" as used herein is to be interpreted accordingly. Generally sequences which are substantially identical will share at least 90%, preferably at least 95% and more preferably at least 98% nucleic acid sequence identity. Sequence identity may be conveniently calculated based on an optimal alignment, for example using the BLAST program accessible at WWW.ncbi.nlm.nih.gov.

The invention will be further understood with reference to the following non-limiting Examples, together with the accompanying Figures in which:

Figure 1 is a plasmid map of the vector pGN1

containing opposable T7 promoters flanking a multiple cloning site and an ampicillin resistance marker.

5 Figure 2 is a plasmid map of the vector pGN8 (a genomic fragment of the *C. elegans* *unc-22* gene cloned in pGN1).

10 Figure 3 is a plasmid map of the vector pGN29 containing two T7 promoters and two T7 terminators flanking *Bst*XI sites. This vector permits cloning of DNA fragments linked to *Bst*XI adaptors.

15 Figure 4 is a plasmid map of the vector pGN39 containing two T7 promoters and two T7 terminators flanking attR recombination sites (based on the Gateway™ cloning system of Life Technologies, Inc).

20 Figure 5 is a plasmid map of the vector pGX22 (a fragment of the *C. elegans* gene C04H5.6 cloned in pGN29).

25 Figure 6 is a plasmid map of the vector pGX52 (a fragment of the *C. elegans* gene K11D9.2b cloned in pGN29).

Figure 7 is a plasmid map of the vector pGX104 (a fragment of the *C. elegans* gene Y57G11C.15 cloned in pGN29).

30 Figure 8 is a plasmid map of the vector pGZ8 (a fragment of the *C. elegans* gene T25G3.2 cloned in pGN39).

35 Figure 9 shows the results of an RNAi experiment in which wild-type (N2) or *nuc-1* strain *C. elegans* in liquid culture were fed with *E. coli* containing the

plasmid pGX22.

Figure 10 shows the results of an RNAi experiment in which wild-type (N2) or *nuc-1* strain *C. elegans* in liquid culture were fed with *E. coli* containing the plasmid pGX52.

Figure 11 shows the results of an RNAi experiment in which wild-type (N2) or *nuc-1* strain *C. elegans* in liquid culture were fed with *E. coli* containing the plasmid pGXGZ8.

Figure 12 shows the results of an RNAi experiment in which wild-type (N2) or *nuc-1* strain *C. elegans* in liquid culture were fed with *E. coli* containing the plasmid pGX104

Example 1

Influence of genetic background on the efficiency of RNAi in *C. elegans*.

5 Introduction

Various different *C. elegans* strains were fed with different bacteria, to test the possibility of RNAi by feeding *C. elegans* with bacteria that produce dsRNA. The possibility of DNA delivery and dsRNA delivery has previously been envisaged by using different bacterial strains. In this experiment the importance of the *C. elegans* strain as receptor of the dsRNA is also shown.

For this experiment the following *E. coli* strains were used:

1. MC1061: F-*araD139* Δ (*ara-leu*)7696 *galE15 galK16* Δ (*lac*)X74 *rpsL* (*Str^r*) *hsdR2* (*r_k⁻ m_k⁺*) *mcrA mcrB1*
- regular host for various plasmids,
- Wertman et al., (1986) Gene 49:253-262,
- Raleigh et al., (1989) in Current Protocols in Molecular Biology eds. Ausubel et al, Publishing associates and Wiley Interscience; New York. Unit 1.4.
2. B21(DE3): F- *ompT(lon)* *hsdS_B* (*r_B⁻ m_B⁻*; an *E. coli* B strain) with DE3, a λ prophage carrying the T7 RNA polymerase gene.
- regular host for IPTG inducible T7 polymerase expression,
- Studier et al. (1990) Meth. Enzymol. 185:60-89
3. HT115 (DE3): F- *mcrA mcrb* IN(*rrnD-rrnE*) 1 λ -*rnc14::tr10* (DE3 lysogen: *lacUV5* promoter-T7polymerase)
- host for IPTG inducible T7 polymerase

expression,

- RNaseIII-,

- Fire A, Carnegie Institution, Baltimore, MD,
Pers. Comm.

5

For this experiment the following *C. elegans* strains
were used:

1. *C. elegans* N2: regular WT laboratory strain
- 10 2. *C. elegans* nuc-1(e1393): *C. elegans* strain with a
reduced endonuclease activity (>95%); condensed
chromatin persists after programmed cell death;
15 ingested (bacterial) DNA in the intestinal lumen
is not degraded. Several alleles are described:
e1392 (strong allele: has been used for the
experiments described below); n887 (resembles
e1392) and n334 (weaker allele)
- Stanfield et al. (1998) East Coast Worm
20 meeting abstract 171,
- Anonymous, Worm Breeder's Gazette 1(1):17b
Hevelone et al. (1988) Biochem. Genet. 26:447-461
- Ellis et al., Worm breeder's Gazette 7(2):44
- Babu, Worm Breeder's gazette 1(2):10
25 - Driscoll, (1996) Brain Pathol. 6:411-425
- Ellis et al., (1991) Genetics 129:79-94

For this experiment the following plasmids were used:

- 30 pGN1: A vector encoding for ampicillin resistance,
harbouring a multiple cloning site between two
convergent T7 promoters.
- pGN8: pGN1 containing a genomic fragment of *unc-22*.
- 35 Decreased *unc-22* expression via RNAi results in a
"twitching" phenotype in *C. elegans*.

Experimental conditions

12-well micro-titer plates were filled with approximately 2 ml of NGM agar per well (1 litre of NGM agar: 15g Agar, 1g peptone, 3g NaCl, 1ml cholesterol solution (5 mg/ml in EtOH), with sterile addition after autoclaving of 9.5 ml 0.1M CaCl₂, 9.5 ml 0.1 ml MgSO₄, 25 ml 1M KH₂PO₄/K₂HPO₄ buffer pH 6 and 5 ml nystatin solution (dissolved 10 mg/ml in 1:1 EtOH:CH₃COONH₄ 7.5 M).

10 The dried plates were spotted with approximately 50 µl of an overnight culture of bacteria. When IPTG induction was required, 50 µl of a 10 mM stock solution of IPTG was dropped on top of the bacteria lawn, and incubated at 37°C for approximately 4 hours. Individual nematodes at the L4 growth stage were then placed in single wells. In each well 4 nematodes, and the plates were further incubated at 20°C for 6 days to allow offspring to be formed. The F1 offspring of the seeded nematodes were tested for the twitching phenotype.

Results

Table 1: Percentage of the offspring that show the twitching phenotype

| | | | |
|----|--------------------|-----------|---------------------|
| 5 | MC1061 | N2 | <i>nuc-1</i> |
| | pGN1 | 0% | 0% |
| | pGN1 + IPTG | 0% | 0% |
| | pGN8 | 0% | 0% |
| | pGN8 + IPTG | 0% | 0% |
| 10 | BL21 (DE3) | | |
| | pGN1 | 0% | 0% |
| | pGN1 + IPTG | 0% | 0% |
| | pGN8 | 20% (+) | >90% (++) |
| | pGN8 + IPTG | 20% (+) | >90% (++±) |
| 15 | HT115 (DE3) | | |
| | pGN1 | 0% | 0% |
| | pGN1 + IPTG | 0% | 0% |
| | pGN8 | 50% (±) | >90% (++) |
| 20 | pGN8 + IPTG | 80% (++) | >90% (+++) |

20 %: percentage twitchers

+ : weak twitching

++ : twitching

+++ : strong twitching

25

Conclusions

The experiment with *E. coli* MC1061 shows that no twitching could be observed in this experiment.

30

Neither the N2 nematodes nor the *nuc-1* nematodes showed any twitchers. This is to be expected as *E. coli* MC1061 does not produce any T7 RNA polymerase, and hence the unc-22 fragment cloned in pGN8 is not

expressed as dsRNA.

5 The experiment with *E. coli* strain BL21(DE3) and
nematode strain N2 shows expected results. BL21(DE3)
harbouring plasmid pGN1 does not result in any
twitching as the pGN1 vector is an empty vector. BL21
(DE3) harbouring PGN8 results in the expression of
unc-22 dsRNA. When this dsRNA is fed to the N2
10 nematode (indirectly by feeding with the bacteria that
produce the dsRNA), this results in a twitching
phenotype, indicating that the dsRNA is able to pass
the gut barrier and is able to perform its interfering
activity.

15 Surprisingly the RNAi effect of the unc-22 dsRNA was
even more pronounced in *C. elegans* strain *nuc-1* than
in the wild type N2 strain. Although one may expect
that the *nuc-1* mutation results in the non-degradation
or at least in a slower degradation of DNA, as the
20 NUC-1 protein is known to be involved in DNase
activity, we clearly observe an enhancement of the
RNAi induced phenotype in *C. elegans* with a *nuc-1*
background. The *nuc-1* mutation has not been cloned
yet, but it has been described that the gene is
25 involved in nuclease activity, and more particularly
DNase activity. If the NUC-1 protein is a nuclease, it
may also have activity on nuclease activity on dsRNA,
which would explain the enhanced RNAi phenotype. The
nuc-1 gene product may be a nuclease, or a regulator
30 of nuclease activity. As the mode of action of RNAi is
still not understood, it is also possible that the
NUC-1 protein is interfering in the mode of action of
RNAi. This would explain why a *nuc-1* mutant is more
sensitive to RNAi.

35

The experiment with the *E. coli* strain HT115 (DE3)

confirms the experiments with the BL21(DE3) strain. The RNA interference observed with the unc-22 dsRNA is even higher. In comparison with strain BL21(DE3) this could be expected, as HT115(DE3) is a RNase III minus strain, and hence is expected to produce larger amounts of dsRNA, resulting in more prominent RNAi. This indicates further that the RNAi observed in this experiment is the result of the dsRNA produced by the bacteria fed to the *C. elegans*. Feeding *C. elegans* nuc-1 with HT115(DE3) harbouring pGN8 also results in higher RNA interference phenotype than feeding the same bacteria to *C. elegans* wild-type strain N2. Once again this indicates that improved RNAi can be realised using a nuclease negative *C. elegans* and more particularly with a with the *C. elegans* nuc-1 (e1392) strain.

Summary

RNA interference can be achieved in *C. elegans* by feeding the worms with bacteria that produce dsRNA. The efficiency of this RNA interference is dependent both on the *E. coli* strain and on the genetic background of the *C. elegans* strain. The higher the level of dsRNA production in the *E. coli*, the more RNAi is observed. This can be realised by using efficient RNA expression systems such as T7 RNA polymerase and RNAase negative strains, such as RNaseIII minus strains. In this example the level of dsRNA production varied: HT115(DE3) > BL21(DE3) > MC1061.

RNA interference is high in *C. elegans* strains that are nuclease negative, or that are influenced in their nuclease activity. This can be realised by using a mutant strain such as *C. elegans* nuc-1. In this example the sensitivity to RNAi varied: *C. elegans* nuc-1 >> *C. elegans* N2

Example 2

Improved RNAi by feeding dsRNA producing bacteria in selected *C. elegans* strains-Comparison of the *nuc-1* strain with several mutants which show improved gut uptake. (designated herein 'gun' mutants). Strains bg77, bg78, bg83, bg84, bg85, bg86, bg87, bg88 and bg89 are typical gun mutant *C. elegans* strains isolated using selection for increased gut uptake (gun phenotype) with the marker dye BCECF-AM.

Experimental conditions:

- 12-well micro-titer plates were filled with approximately 2ml of NGM agar (containing 1ml/l of ampicillin (100µg/ml) and 5 ml of 100mM stock IPTG) per well
- the dried plates were spotted with 25µl of an overnight culture of bacteria (BL21DE3/HT115DE3) containing the plasmids pGN1 (T7prom-T7prom) or pGN8 (T7prom-unc-22-T7prom)
- individual nematodes at the L4 growth stage were then placed in single wells, one nematode per well
- the plates were incubated at 20°C for 6 days to allow offspring to be formed
- the adult F1 offspring of the seeded nematodes were tested for the twitching phenotype

Results:**Table 2:**

| | 20°C/6d | pGN1 HT115DE3 | pGN8 BL2DE3 | pGN8 HT115DE3 | |
|----|---------|------------------|-------------|------------------|--|
| 5 | N2 | 0 | 1 | 1 | |
| | bg77 | 0 | 1-2 | 3 | |
| | bg78 | 0 | 1 | 1-2 | |
| | bg83 | 0 | 1 | 1 | |
| 10 | bg84 | 0 | 1-2 | 3 | |
| | bg85 | 0 | 1 | 2-3 | |
| | bg86 | 0 | 1 | 2 | |
| | bg87 | 0 | 1 | 1 | |
| | bg88 | 0 | 1 | 1 | |
| 15 | bg89 | 0 | 1 | 1 | |
| | | | | | |

figure legend:

- 0 = no twitching
 1 = no to weak phenotype
 2 = clear phenotype
 3 = strong phenotype

Conclusions

- bacterial strain HT115(DE3) shows a better RNAi sensitivity than bacterial strain BL21(DE3)
- the *nuc-1* *C. elegans* strain is a better strain than the Wild-type N2 strain for RNAi sensitivity
- various gun mutants (improved gut uptake mutants) and more particularly the gun mutant strains bg77, bg84, bg85, bg86 show improved sensitivity to RNAi compared to Wild-type.

- 22 -

A double mutant *C. elegans* strain (nuc-1/gun) shows even greater sensitivity to RNAi compared to wild-type:

5 Double mutants were constructed to test the prediction that gun/nuc mutants would even show more enhanced RNAi sensitivity. As an example, the crossing strategy with gun strain bg85 is shown, similar crosses can be conducted with other gun strains, such
10 as bg77, bg84 and bg86.

P0 cross: gun(bg85) x WT males

15 F1 cross: nuc-1 x gun(bg85)/+ males

F2 cross: nuc-1 x gun(bg85)/+; nuc-1/0 males (50%)
nuc-1 x +/+; nuc-1/0 males (50%)

20 F3 single: gun(bg85)/+; nuc-1 hermaphrodites (25%)
+/+; nuc-1 hermaphrodites (75%)

F4 single: gun(bg85); nuc-1 (1/4 of every 4th plate high staining with BCECF)

25 F5 retest: gun(bg85); nuc-1 (100% progeny of F4 singled high staining with BCECF)

To select for the gun phenotype, the fluorescence precursor BCECF-AM is used (obtainable from Molecular probes). The precursor BCECF-AM is cleaved by
30 esterases present in the gut of the worm to generate the dye BCECF which is fluorescent at pH values above 6. This allows selection for worms that have a gun phenotype. BCECF-AM is taken up through the pharynx
35 into the gut lumen and is not fluorescent until it has been cleaved, and the BCECF portion has entered the

cells surrounding the lumen. Wild-type worms will show slower or no increase in BCECF fluorescence.

5 **Example 3**

Improved RNAi feeding in liquid culture using *nuc-1*(e1393) *C. elegans*.

Introduction

10 N2 and *nuc-1* *C. elegans* strains were fed with bacteria producing dsRNAs that give lethal phenotypes via RNAi. For this example RNAi was performed in liquid culture instead of on agar plates. We show here for a number of genes that the RNAi effect is more penetrant using
15 the *nuc-1* strain than the N2 strain, and that RNAi can be performed in liquid.

For this experiment the following *E. coli* strains were used:

20

1. HT115 (DE3): F- *mcrA mcrB* IN(*rrnD-rrnE*) 1 λ -*rncl4::trl0* (DE3 lysogen: lacUV5 promoter -T7 polymerase)
 - host for IPTG inducible T7 polymerase expression
 - 25 - RNaseIII⁻
 - Fire A, Carnegie Institution, Baltimore, MD, Pers. Comm.

25

For this experiment, following *C. elegans* strains were used:

30

1. *C. elegans* N2: regular WT laboratory strain
2. *C. elegans nuc-1*(e1393): *C. elegans* strain with a
35 reduced endonuclease activity (>95%); condensed chromatin persists after programmed cell death;

35

- ingested (bacterial) DNA in the intestinal lumen is not degraded. Several alleles are described: e1392 (strong allele: has been used for the experiments described below); n887 (resembles e1392) and n334 (weaker allele)
- 5 - Stanfield et al. (1998) East Coast Worm meeting abstract 171
- Anonymous, Worm Breeder's Gazette 1(1):17b
- Hevelone et al. (1988) Biochem. Genet. 26:447-461
- 10 - Ellis et al., Worm breeder's Gazette 7(2):44
- Babu, Worm Breeder's gazette 1(2):10
- Driscoll, (1996) Brain Pathol. 6:411-425
- Ellis et al., (1991) Genetics 129:79-94

15

For this experiment, the following plasmids that all give lethal phenotypes in *C. elegans* via RNAi were used:

- 20 pGX22: a vector encoding ampicillin resistance, containing a genomic fragment of cosmid C04H5.6 corresponding to a member of the RNA helicase family.

- pGX52: a vector encoding ampicillin resistance, containing a genomic fragment of cosmid K11D9.2b corresponding to sarco/endoplasmic Ca²⁺ ATPase also known as SERCA.
- 25

- pGZ18: a vector encoding ampicillin resistance, containing a genomic fragment of cosmid T25G3.2 corresponding to a chitin like synthase gene.
- 30

- pGX104: a vector encoding ampicillin resistance, containing a genomic fragment of cosmid Y57G11C.15 corresponding to sec-61, a transport protein.
- 35

Experimental conditions

- 5 - 1 ml overnight cultures of HT115 (DE3) bacteria containing the plasmids pGX22, pGX52, pGZ18 or pGX104 respectively were pelleted and resuspended in S-complete medium, containing 1ml/l of ampicillin (100 µg/ml) and 1ml/l of 1000mM IPTG.
- 10 - 10 µl of this bacterial solution was transferred to a 96-well microtiter plate already filled with 100 µl S-complete containing 1ml/l of ampicillin (100 µg/ml) and 1ml/l of 1000mM IPTG.
- 15 - 3 nematodes at the L1 growth stage of N2 and nuc-1 strain were then placed in single wells, 3 L1's per well. Per experimental set up, 16 wells were used (n=16).
- 20 - the plates were incubated at 25°C for 5 days to allow offspring to be formed.
- 25 - the plates were visually checked and the following phenotypes could be scored per individual well:
 - 25 **no effect:** L1's developed to adults and gave normal offspring.
 - 30 **no F1 offspring:** L1's developed to adults and gave no offspring.
 - 30 **acute lethal:** original L1 did not mature and died.

Results

- 35 The results of this experiment are illustrated graphically in Figures 9 to 12. Data are expressed as

a percentage of the total (n=16) on the y-axis for both N2 and nuc-1 strains.

Conclusions

5 The following genes were tested in this liquid RNAi assay:

- C04H5.6: an RNA helicase. RNAi of this gene interferes with the generation of offspring.
- 10 - SERCA: a sarco/endoplasmic Ca^{2+} ATPase. A strong RNAi phenotype causes an acute lethal phenotype. A less penetrant RNAi effect results in loss of offspring.
- T25G3.2: a chitin like synthase gene. RNAi of
15 this gene causes dead eggs.
- sec-61: a transport protein. A strong RNAi phenotype causes an acute lethal phenotype. A less penetrant RNAi effect results in loss of offspring.
- 20 - RNAi can be performed under liquid conditions.

As in the previous examples this set of experiments shows that the nuc-1 *C. elegans* strain is more sensitive to RNAi than the wild-type N2 strain. This
25 is most clear for less penetrant phenotypes such as SERCA and chitin synthase. For strong RNAi phenotypes like the helicase and Sec-61 the difference between the N2 wild-type strain and the nuc-1 stain is less pronounced.

30

Example 4**Cloning of pGX22, pGX52, pGZ18 and pGX104 for RNAi**

A set of primers for each gene was designed on the basis of sequence data available in the publicly accessible *C. elegans* sequence database (Acedb).

The cosmid names relate to:

1. **C04H5.6**=member of RNA helicase
2. **K11D9.2b**=SERCA
3. **Y57G11C.15**=transport protein sec-61
4. **T25G3.2**=chitin synthase like

The following primer sequences were designed:

1. **C04H5.6F** 5'-TGCTCAGAGAGTTTCTCAACGAACC-3'
C04H5.6R 5'-CAATGTTAGTTGCTAGGACCACTG-3'
2. **K11D9.2bF** 5'-CAGCCGATCTCCGTCTTGTG-3'
K11D9.2bR 5'-CCGAGGGCAAGACAACGAAG-3'
3. **Y57G11C.15F** 5'-ACCGTGGTACTCTTATGGAGCTCG-3'
Y57G11C.15R 5'-TGCAGTGGATTGGGTCTTCG-3'
4. **T25G3.2F**
5'-GGGGACAAGTTTGTACAAAAAGCAGGCTATGCCAAGTACATGTCGATTGCG-3'
T25G3.2R
5'-GGGGACCAC'TTTGTACAAGAAAGCTGGGTGGAGAAGCATTCGAGAGTTTG-3'

PCR was performed on genomic DNA of N2 strain *C. elegans* to give PCR products of the following sizes:

- 1326bp for C04H5.6
- 1213bp for K11D9.2b

1024bp for Y57G11C.15

1115bp for T25G3.2

- 5 The PCR fragments of C04H5.6, K11D9.2b and Y57G11C.15 were linked to *Bst*XI adaptors (Invitrogen) and then cloned into the pGN29 vector cut with *Bst*XI. pGN29 contains two T7 promoters and two T7 terminators flanking a cloning site which is adapted for facilitated cloning of PCR fragments, comprising a
- 10 stuffer DNA flanked by two *Bst*XI sites (see schematic Figure 3). The resulting plasmids were designated pGX22 (C04H5.6), pGX52 (K11D9.2b) and pGX104 (Y57G11C.15).
- 15 The PCR fragment of T25G3.2 was cloned into pGN39 via recombination sites based on the GATEWAY™ cloning system (Life Technologies, Inc). pGN39 contains two T7 promoters and two T7 terminators flanking a cloning site which facilitates "High Throughput" cloning based
- 20 on homologous recombination rather than restriction enzyme digestion and ligation. As shown schematically in Figure 4, the cloning site comprises attR1 and attR2 recombination sites from bacteriophage lambda flanking a gene which is lethal to *E. coli*, in this
- 25 case the *ccdB* gene. This cloning site is derived from the Gateway™ cloning system commercially available from Life Technologies, Inc. The Gateway™ cloning system has been extensively described by Hartley et al. in WO 96/40724 (PCT/US96/10082).

Example 5

Selecting *C. elegans* mutations for increased gut uptake (gun) using marker dye BCECF-AM and *unc-31* as background.

5

The screen was performed in *unc-31(e928)* mutant background, to ensure high amounts of dye in the gut lumen, since *unc-31* mutations show constitutive pharyngeal pumping. The dye (BCECF-AM: 2',7' bis (2-carboxyethyl)-5-(and-6)-carboxyfluorescein, acetoxymethylester), obtained from Molecular Probes, is cleaved by intracellular esterases. Fluorescence accumulates in the gut cells upon passage through the apical gut membrane.

15

Mutagenesis

Day 1: *unc-31* L4 staged worms were mutagenised with EMS (final concentration 50 mM) for 4 hours

Day 2: P0 was divided over several large agar plates

Day 6: F1's were collected and dropped on large plates. The number of eggs the F1's layed were checked every hour and de F1's were removed when 10-20 eggs per F1 were counted

Day 10: F2 adults were collected and screened with BCECF-AM. Mutations with increased staining of the gut cells after 15-30 minutes exposure to the dye were selected and singled on small agar plates.

30

About 50 initial positives gave progeny which was retested with BCECF-AM (2x) and leucine CMB (1x) 9 of the 50 strains were kept (2 strains : 3 times positive, 7 other strains : twice positive)

35

Table 3: Isolation of mutations for increased staining with BCECF-AM

| Total P0 | Total F1 | Total F2 | screened chromosomes | number of strains isolated |
|-----------|-------------|--------------|----------------------|----------------------------|
| (counted) | (estimated) | (calculated) | (estimated) | (counted) |
| 2251 | 55618 | 222472 | 100000 | 9 |

Outcrossing, backcrossing and double construction

- 10 1. backcrossing *unc-31; gun* --> *unc-31; gun*
 - *unc-31; gun* x WT males
 - singled 2x5 WT hermaphrodites F1s (*unc-31/+; gun/+*)
 - singled 50 WT hermaphrodites F2s (1/4 homozygous)
 - select strains segregating 1/4 *unc*
- 15 - stain *unc* strains with BCECF-AM
 - from positive strains pick *unc* homozygous
 - retest 100 % *unc* strains with BCECF-AM
 - kept 1 strain (backcrossed)
- 20 2. *unc-31* background was crossed out-->+; *gun*
 - *unc-31; gun* x WT males
 - singled 2x5 WT hermaphrodites F1s (*unc-31/+; gun/+*)
 - singled 50 WT hermaphrodites F2s (1/4 homozygous)
 - select strains which did not segregate *unc* F3s
- 25 anymore
 - stain non *unc* strains with BCECF-AM
 - 7 positive strains were retested with BCECF-AM and finally 1 was selected and kept (outcrossed)
- 30 3. +; *gun* (1x outcrossed) were 2 times backcrossed-->+; *gun* (3x backcrossed)
 - *gun* x WT males
 - WT hermaphrodites x F1 males (*gun/+*)
 - singled 10 WT hermaphrodites F2s (1/2 heterozygous)
- 35 - singled 50 WT hermaphrodites F3s (1/8 homozygous)

- 31 -

- stain strains with BCECF-AM- retested positives with BCECF-AM and finally 1 was selected and kept

4. *gun* (3x backcrossed) were crossed with *nuc-1(X)*
 5 mutant--> *gun; nuc-1*
 - *gun* x WT males
 - *nuc-1* x *gun/+* males
 - *nuc-1* x *gun/+; nuc-1/0* or *+/+; nuc-1/0* males
 - singled 10 WT hermaphrodite progeny (*nuc-1*
 10 homozygous, $\frac{1}{2}$ heterozygous *gun*)
 - singled 40 WT hermaphrodite progeny (1/8 homozygous *gun*)
 - stain strains with BCECF-AM
 - retested positives with BCECF-AM and finally 1 was
 15 selected and kept

Table 6: Strains derived from *gun* mutations

| | <i>gun</i> | <i>unc-31; gun</i> | | <i>unc-31; gun</i> | | <i>+/; gun</i> | | | <i>gun; nuc-1</i> |
|----|------------------|--------------------|---------------|--------------------|---------------|------------------|---------------|---------------|-------------------|
| | original isolate | backcrossed (1x) | | outcrossed (1x) | | 3x b.c. | | from 3x b.c. | |
| | allele number | isolation number | strain number | isolation number | strain number | isolation number | strain number | strain number | strain number |
| 20 | bg77 | 31.4 | UG 510 | 31.4.46.1 | UG 556 | 31.4.34 | UG 563 | UG 674 | UG 777 |
| 25 | bg78 | 37.5 | UG 511 | 37.5.46.4 | UG 557 | 37.5.15 | UG 564 | UG 675 | - |
| | bg83 | 10.2 | UG 543 | 10.2.11 | UG 600 | 10.2.21 | UG 586 | UG 676 | - |
| | bg84 | 7.2 | UG 544 | 7.2.10 | UG 601 | 7.2.15 | UG 589 | UG 677 | UG 774 |
| | bg85 | 11.5 | UG 545 | 11.5.29.2 | UG 602 | 2x b.c. | UG 717 | | UG 775 |
| | bg86 | 42.1 | UG 546 | 42.1.4.5 | UG 603 | 42.1.18 | UG 587 | UG 678 | UG 776 |
| 30 | bg87 | 7.1 | UG 547 | 7.1.8.3 | UG 604 | 7.1.22 | UG 585 | UG 679 | - |
| | bg88 | 5.3 | UG 548 | 5.3.9 | UG 605 | 5.3.18 | UG 584 | UG 680 | - |
| | bg89 | 23.4 | UG 549 | 23.4.13.5 | UG 606 | 23.4.3 | UG 588 | UG 671 | - |

35

SEQUENCE LISTING:

SEQ ID NO: 1 complete sequence of pGN1

5 SEQ ID NO: 2 complete sequence of pGN8

SEQ ID NO: 3 complete sequence of pGN29

SEQ ID NO: 4 complete sequence of pGN39

10 SEQ ID NO: 5 complete sequence of pGX22

SEQ ID NO: 6 complete sequence of pGX52

15 SEQ ID NO: 7 complete sequence of pGX104

SEQ ID NO: 8 complete sequence of pGZ8

SEQ ID NO: 9 primer C04H5.6F

20 SEQ ID NO: 10 primer C04H5.6R

SEQ ID NO: 11 primer K11D9.2bF

25 SEQ ID NO: 12 primer K11D9.2bR

SEQ ID NO: 13 primer Y57G11C.15F

SEQ ID NO: 14 primer Y57G11C.15R

30 SEQ ID NO: 15 primer T25G3.2F

SEQ ID NO: 16 primer T25G3.2R

Claims:

1. A method of inhibiting expression of a target gene in a nematode worm comprising feeding to
5 said nematode worm a food organism which is capable of producing a double-stranded RNA structure having a nucleotide sequence substantially identical to a portion of said target gene following ingestion of the food organism by the nematode, wherein the nematode
10 has a non wild-type genetic background selected to provide increased sensitivity to RNA interference as compared to wild type.
2. A method as claimed in claim 1 wherein the
15 nematode is a microscopic nematode.
3. A method as claimed in claim 2 wherein the nematode is from the genus *Caenorhabditis*.
- 20 4. A method as claimed in claim 3 wherein the nematode is *C. elegans*.
5. A method as claimed in any one of claims 1 to 4 wherein the nematode has a mutant genetic
25 background.
6. A method as claimed in claim 5 wherein the nematode is a mutant strain which exhibits reduced activity of one or more nucleases compared to wild
30 type.
7. A method as claimed in claim 6 wherein the nematode is *C. elegans* strain *nuc-1*.
- 35 8. A method as claimed in claim 5 wherein the nematode is a mutant strain which exhibits increased

gut uptake compared to wild type.

9. A method as claimed in claim 8 wherein the nematode is mutant *C. elegans* strain bg85.

5

10. A method as claimed in claim 5 wherein the nematode is a mutant strain having at least one mutation which results in reduced nuclease activity compared to wild type and at least one mutation which results in increased gut uptake compared to wild type.

10

11. A method as claimed in claim 10 wherein the nematode is a mutant *C. elegans* strain having the *nuc-1* mutation and the bg85 mutation.

15

12. A method as claimed in any one of the preceding claims wherein the food organism has been engineered to express a double-stranded RNA.

20

13. A method as claimed in any one of the preceding claims wherein the food organism is a bacterium.

25

14. A method as claimed in claim 13 wherein the food organism is *E. coli*.

15. A method as claimed in any one of the preceding claims wherein the food organism has been genetically modified to express a double-stranded RNA having a nucleotide sequence substantially identical to a portion of said target gene.

30

16. A method as claimed in claim 15 wherein the food organism contains a DNA vector, the vector comprising a promoter or promoters orientated relative to a DNA sequence such that they are capable of

35

initiating transcription of said DNA sequence to RNA capable of forming a double-stranded structure upon binding of an appropriate RNA polymerase to said promoter or promoters.

5

17. A method as claimed in claim 25 wherein the vector comprises two promoters flanking the DNA sequence.

10

18. A method as claimed in claim 26 wherein the two promoters are identical.

15

19. A method as claimed in claim 25 wherein the vector comprises a single promoter and further comprises said DNA sequence in a sense and an antisense orientation relative to said promoter.

20

20. A method as claimed in any one of claims 16 to 20 wherein the nematode or the food organism is adapted to express an RNA polymerase capable of initiating transcription from said promoter or promoters.

25

21. A method as claimed in any one of claims 16 to 20 wherein the RNA polymerase is T7, T3 or SP6 polymerase.

30

22. A method as claimed in any one of claims 1 to 21 wherein the step of feeding said food organism to said nematode worm is carried out in liquid culture.

35

23. A method of inhibiting expression of a target gene in a nematode worm comprising feeding to said nematode worm a food organism capable of producing a double-stranded RNA structure having a nucleotide sequence substantially identical to a

portion of said target gene following ingestion of the food organism by the nematode, wherein the food organism carries a modification selected to provide increased expression or persistence of the doubled-stranded RNA compared to a food organism which does not carry the modification.

24. A method as claimed in claim 23 wherein the food organism is a bacterium.

25. A method as claimed in claim 24 wherein the bacterium is an *E. coli* strain.

26. A method as claimed in claim 25 wherein the *E. coli* strain is an RNase III minus strain or any other RNase negative strain.

27. A method as claimed in any one of claims 23 to 26 wherein the step of feeding said food organism to said nematode worm is carried out in liquid culture.

28. A method of inhibiting expression of a target gene in a nematode worm comprising introduction of a DNA capable of producing a double-stranded RNA structure having a nucleotide sequence substantially identical to a portion of said target gene in said nematode, wherein the nematode is one which exhibits increased gut uptake compared to wild type.

29. A method as claimed in claim 28 wherein the nematode is a microscopic nematode.

30. A method as claimed in claim 29 wherein the nematode is from the genus *Caenorhabditis*.

31. A method as claimed in claim 30 wherein the

nematode is *C. elegans*.

32. A method as claimed in any one of claims 28
to 31 wherein the nematode has a mutant genetic
5 background.

33. A method as claimed in claim 32 wherein the
nematode is mutant *C. elegans* strain bg85.

10 34. A method as claimed in any one of claims 28
to 33 wherein the DNA capable of producing a double-
stranded RNA structure is a vector comprising a
promoter or promoters orientated relative to a DNA
sequence such that they are capable of initiating
15 transcription of said DNA sequence to RNA capable of
forming a double-stranded structure upon binding of an
appropriate RNA polymerase to said promoter or
promoters.

20 35. A method as claimed in claim 34 wherein the
vector comprises two promoters flanking the DNA
sequence.

25 36. A method as claimed in claim 35 wherein the
two promoters are identical.

30 37. A method as claimed in claim 34 wherein the
vector comprises a single promoter and further
comprises said DNA sequence in a sense and an
antisense orientation relative to said promoter.

35 38. A method as claimed in any one of claims 34
to 37 wherein the nematode is adapted to express an
RNA polymerase capable of initiating transcription
from said promoter or promoters.

39. A method as claimed in any one of claims 34

to 38 wherein the RNA polymerase is T7, T3 or SP6 polymerase.

FIG. 1




FIG. 2.




FIG. 3.




FIG. 4.




FIG. 5.




FIG. 6.




FIG. 7.




FIG. 8.




FIG. 9.




FIG. 10.




FIG. 11.





FIG. 12.



1
SEQUENCE LISTING

<110> DEVGEN NV

<120> IMPROVEMENTS RELATING TO DOUBLE-STRANDED RNA INHIBITION

<130> SCB/53711/001

<140>

<141>

<160> 14

<170> PatentIn Ver. 2.0

<210> 1

<211> 3216

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Plasmid pGN1

<400> 1

| | | | | | | |
|------------|------------|------------|------------|-------------|------------|------|
| gagtgcacca | tatgcggtgt | gaaataccgc | acagatgcgt | aaggagaaaa | taccgcatca | 60 |
| ggcgaaattg | taaacgttaa | tattttggtt | aaattcgcgt | taaatatttg | ttaaatcagc | 120 |
| tcatttttta | accaataggc | cgaaatcggc | aaaatccctt | ataaatcaaa | agaatagacc | 180 |
| gagataggg | tgagtgttgt | tccagtttgg | aacaagagtc | cactattaaa | gaacgtggac | 240 |
| tccaacgtca | aagggcgaaa | aaccgtctat | cagggcgatg | gccactacg | tgaaccatca | 300 |
| cccaaatcaa | gttttttgcg | gtcaggtgtc | cgtaaagctc | taaatcggaa | ccctaaaggg | 360 |
| agccccgat | ttagagcttg | acggggaaa | ccggcgaa | tgccgagaaa | ggaagggag | 420 |
| aaagcgaaa | gagcgggcgc | tagggcgctg | gcaagtgtag | cggtcacgct | gcgcgtaacc | 480 |
| accacaccgc | ccgcgcttaa | tgccgcgcta | cagggcgctg | ccattcgcca | ttcaggctgc | 540 |
| gcaactgttg | ggaagggcga | tcggtgcggg | cctcttcgct | attacgccag | ctggcgaaa | 600 |
| gggatgtgc | tgcaaggcga | ttaagttggg | taacgccagg | gttttcccag | tcacgacgtt | 660 |
| gtaaaacgac | ggccagtga | ttgtaatacg | actcactata | gggcgaattc | gagctcggtt | 720 |
| cccggggatc | ctctagagtc | gaaagcttct | cgccctatag | tgagtcgtat | tacagcttga | 780 |
| gtattctata | gtgtcaccta | aatagcttgg | cgtaatcatg | gtcatagctg | tttcctgtgt | 840 |
| gaaattgtta | tccgctcaca | attccacaca | acatacgagc | cggaagcata | aagtgtaaa | 900 |
| cctggggtgc | ctaatagtgt | agctaactca | cattaattgc | gttgcgctca | ctgcccgctt | 960 |
| tccagtcggg | aaacctgtcg | tgccagctgc | attaatgaat | cgcccaacgc | gcggggagag | 1020 |
| gcggtttg | tattgggcgc | tcttcgctt | cctcgctcac | tgactcgctg | cgctcggtcg | 1080 |
| ttcggtcg | gcgagcggt | tcagctcact | caaaggcggt | aatacggtta | tccacagaat | 1140 |
| caggggataa | cgaggaaa | aacatgtgag | caaaaggcca | gcaaaaggcc | aggaaccgta | 1200 |
| aaaaggccgc | gttgctggcg | tttttcgata | ggctccgccc | ccctgacgag | catcacaata | 1260 |
| atcgacgtc | aagtcagagg | tgccgaaa | cgacaggact | ataaagatac | caggcggttc | 1320 |
| cccctggaag | ctccctcg | cgctctcctg | ttccgaccct | gcccgttacc | ggatacctgt | 1380 |
| ccgcctttct | cccttcggga | agcgtggcgc | tttctcatag | ctcacgctgt | aggtatctca | 1440 |
| gttcggtgta | ggtcggttcg | tccaagctgg | gctgtgtgca | cgaaccccc | gttcagccc | 1500 |
| accgctgcgc | cttatccggt | aactatcg | ttgagtc | cccggttaaga | cacgacttat | 1560 |
| cgccactggc | agcagccact | ggtaacagga | ttagcagagc | gaggtatgta | ggcggtgcta | 1620 |
| cagagttctt | gaagtgtgg | cctaactacg | gtacactag | aaggacagta | tttggtatct | 1680 |
| gcgctctgct | gaagccagtt | accttcggaa | aaagagttgg | tagctcttga | tccggcaaac | 1740 |
| aaaccaccgc | tggtagcgg | ggtttttttg | tttgcaagca | gcagattacg | cgcaaaaaa | 1800 |
| aaggatctca | agaagatcct | ttgatctttt | ctacggggtc | tgacgctcag | tggaacgaaa | 1860 |
| actcacgtta | agggattttg | gtcatgagat | tatcaaaaag | gatcttcacc | tagatccttt | 1920 |
| taaattaaaa | atgaagtttt | aatcaatct | aaagtatata | tgagtaaact | tggtctgaca | 1980 |

```

gttaccaatg cttaatcagt gaggcaccta tctcagcgat ctgtctatctt cgttcatcca 2040
tagttgcctg actccccgtc gtgtagataa ctacgatacg ggagggccta ccatctggcc 2100
ccagtgcctg aatgataccg cgagacccac gctcacccgc tccagattta tcagcaataa 2160
accagccagc cgggaagggcc gagcgagaa gtggtcctgc aactttatcc gcctccatcc 2220
agtctattaa ttgttgccgg gaagctagag taagtagttc gccagttaat agtttgcgca 2280
acgttgcttg cattgctaca ggcatcgtgg tgtcacgctc gtcgtttggt atggcttcat 2340
tcagctccgg ttcccaacga tcaaggcgag ttacatgac ccccatgttg tgcaaaaaag 2400
cggttagctc cttcggctcct ccgatcgttg tcagaagtaa gttggccgca gtgttatcac 2460
tcattggtat ggcagcactg cataattctc ttactgtcat gccatccgta agatgctttt 2520
ctgtgactgg tgagtactca accaagtcat tctgagaata ccgcgcccg cgaccgagtt 2580
gctcttgccc ggcgtcaata cgggataata gtgtatgaca tagcagaact ttaaaagtgc 2640
tcattcattg aaaacgttct cggggcgcaa aactctcaag gatcttaccg ctgttgagat 2700
ccagttcgat gtaaccact cgtgcaccca actgatcttc agcatctttt actttcacca 2760
gcgtttctgg gtgagcaaaa acaggaaggc aaaatgccgc aaaaaaggga ataagggcga 2820
cacggaaatg ttgaatactc atactcttcc tttttcaata ttattgaagc atttatcagg 2880
gttattgtct catgagcggg tacatatttg aatgtattta gaaaaataaa caaatagggg 2940
ttccgcgcac atttcccga aaagtgcac ctgacgtcta agaaaccatt attatcatga 3000
cattaacctc taaaaatagg cgtatcacga ggcccttcg tctcgcgcgt ttcggtgatg 3060
acggtgaaaa cctctgacac atgcagctcc cggagacggt cacagcttgt ctgtaagcgg 3120
atgccgggag cagacaagcc cgtcagggcg cgtcagcggg tgttggcggg tgtcggggct 3180
ggcttaacta tgcggcatca gaggcagatt tactga 3216

```

<210> 2

<211> 4620

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Plasmid pGN8

<400> 2

```

gatccgaatc tccatgtctg ttaacagcct tgacacggaa tttatattca tgccottgag 60
tcaaactcgtc aacgtggaag ttggtatcct tgctctctcc gcaagcagtc catctgccag 120
tggcagcatc ttgcttttca atgacatagt gactgatttc agctcctcca tcatctctctg 180
gttccttcca tgcaagatca catccatcct tgacaatatt agtgacatcg agagggtccac 240
gtgggcttga tggatgatca agaacagtaa ccttcacttc agcagtgtca gttccattct 300
cgttctctgc cttgatgata taggttcctg tatccgaacg caaagctctc ttcacatgga 360
atttagtctt gccgtcttca ttgttcaact tcatacagtc atcagattcg actggtgttc 420
cttcgaaagt ccaagtaatt gtggagttg gttcaccact gactggaatg ttcaatgaga 480
agtcttgctc agccttgacc ttgatttctt gaatcgagtt acgatcgatg actgggtggaa 540
ctataattta attcaatgat tattagtaat tgatttagac tcttaccatt tctagccttt 600
gcaacagctg atgctgaatc agatggatct cccaatcctg ccttggtctt ggcacggatt 660
ctgaattcgt actttgatcc ttcttgaga tttccaacag tagcattcgt ttgtccagct 720
ggaacatgag caacgtcatt ccagaatggc gagaactcgt ccttcacttc aacaacgtat 780
tcctcgattg gagcaccacc gtcggttgct ggtggcttcc attcaagtc aacatgatcc 840
ttatcccaat cagtaatttc aggagcattt gtctttcctg gcttgctaaa tggatctttg 900
gcaagtgtgg ttccgaaggt ctccaatgga tcggactctc cttcagcatt gacggcagcg 960
acacggaact gaaaatcaaa atgttgtagg caattgagtt caagattaaa aaattctcac 1020
tttatattca tgtccaggaa taagaccgtc aacaacagct gtagtcttat ctccagcgac 1080
ctttgcagct ggaaccatc ttccacttgc agtatcgtac ttttcgatca catagttttc 1140
aattggaata cctccatcat catctggtgc acgccaatc aaagtgcac gatcaccatg 1200
aacatcggaa acatctaatt gaccatttgg agaagtggc ttgtctgaaa atttaaaata 1260
taaccaaat aattgaagaa aactaatgct caccaataac attgatctta acagttgctt 1320
catcttctcc atttgcatg acagcttga tagtgaaagt tccactgtct ccacgttcca 1380
tttgcttca aaccagcttt gattgggtatt ctgggttatc aagcttctcg ccttatagtg 1440
agtcgtatta cagcttgagt attctatagt gtcacctaaa tagcttgcg taatcatggt 1500
catagctgtt tcctgtgtga aattgttate cgctcacaat tccacacaac atacgagcgg 1560
gaagcataaa gtgtaaagcc tggggtgcct aatgagttag ctaactcaca ttaattgcgt 1620

```

```

tgcgctcact gcccgctttc cagtcgggaa acctgtcgtg ccagctgcat taatgaatcg 1680
gccaacgcgc ggggagaggc gggttgcgta ttgggcgctc ttccgcttcc tcgctcactg 1740
actcgctgcg ctccggtcggt cggctgcggc gagcgggtatc agctcactca aaggcggtaa 1800
tacggttatc cacagaatca ggggataacg caggaaagaa catgtgagca aaaggccagc 1860
aaaaggccag gaaccgtaaa aaggccgcgt tgctggcggt ttccgatagg ctccgcccc 1920
ctgacgagca tcacaaaaat cgacgctcaa gtcagagggt gcgaaacccg acaggactat 1980
aaagatacca ggcgtttccc cctggaagct ccctcgtgcg ctctcctgtt ccgaccctgc 2040
cgcttaccgg atacctgtcc gcctttctcc ctccgggaag cgtggcgctt tctcatagct 2100
cacgctgtag gtatctcagt tcggtgtagg tcgttcgctc caagctgggc tgtgtgcacg 2160
aacccccgt tcagcccgac cgctgcgcct tatccggtaa ctatcgtctt gagtccaacc 2220
cggtaagaca cgacttatcg ccactggcag cagccactgg taacaggatt agcagagcga 2280
ggatgtgtag cgggtgctaca gagtcttga agtgggtggc taactacggc tacactagaa 2340
ggacagtatt tggatatctgc gctctgctga agccagttac ctccgaaaaa agagttggta 2400
gctcttgatc cggcaaacaa accaccgctg gtagcgggtg tttttttgtt tgcaagcagc 2460
agattacgcg cagaaaaaaa ggatctcaag aagatccctt gatcttttct acggggtctg 2520
acgctcagtg gaacgaaaaa tcacgttaag ggattttgtt catgagatta tcaaaaagga 2580
tcttacccta gatcctttta aattaaaaat gaagttttta atcaatctaa agtatatatg 2640
agtaaacctg gctgacagt taccaatgct taatcagtga ggcacctatc tcagcgatct 2700
gtctatttctg ttcattccata gttgcctgac tcccgcgtcg ttagataact acgatacggg 2760
agggcttacc atctggcccc agtgctgcaa tgataccgcg agaccacgc tcaccggctc 2820
cagatttatc agcaataaac cagccagccg gaagggccga gcgcagaagt ggtcctgcaa 2880
ctttatccgc ctccatccag tctattaatt gttgccggga agctagagta agtagttcgc 2940
cagtttaatac tttgcgcaac gttgttgga gttgtacagg catcgtggtg tcacgctcgt 3000
cgtttggtat ggcttcatc agctccggtt cccaacgatc aaggcgagtt acatgatccc 3060
ccatgttgtg caaaaaagcg gttagctcct tcggtcctcc gatcgttgtc agaagtaagt 3120
tgggcgcagt gttatcactc atggttatgg cagcactgca taattctctt actgtcatgc 3180
catccgtaag atgcttttct gtgactgggt agtactcaac caagtcattc tgagaatacc 3240
gcgcccggcg accgagttgc tcttggcccg cgtcaatacg ggataatagt gtatgacata 3300
gcgaactttt aaaagtgtc atcattggaa aacgttcttc ggggcgaaaa ctctcaagga 3360
tcttaccgct gttgagatcc agttcgatgt aaccactcgc tgcacccaac tgatcttcag 3420
catcttttac tttcaccagc gtttctgggt gagcaaaaac aggaaggcaa aatgccgcaa 3480
aaaagggaat aaggcgacaa cggaatggt gaatactcat actcttcctt tttcaatatt 3540
attgaagcat ttatcagggt tattgtctca tgagcggata catatttgaa tgtatttaga 3600
aaaaataaca aatagggtt ccgcgcacat tccccgaaa agtgccacct gacgtctaag 3660
aaaccattat tatcatgaca ttaacctata aaaataggcg tatcacgagg cccttctcgtc 3720
tcgcgcgttt cgggtgatgac ggtgaaaacc tctgacacat gcagctcccg gagacggta 3780
cagcttgtct gtaagcggat gccgggagca gacaagccc tcagggcgcg tcagcgggtg 3840
ttggcgggtg tcggggctgg cttaactatg cggcatcaga gcagattgta ctgagagtgc 3900
accatattgc gtgtgaaata ccgcacagat gcgtaaggag aaaataaccgc atcaggcgaa 3960
attgtaaacg ttaatatttt gttaaaatt gcgttaataa tttgttaaat cagctcattt 4020
tttaaccaat aggcggaaat cggcaaaatc ccttataaat caaaagaata gaccgagata 4080
gggttgagtg ttgttccagt ttggaacaag agtccactat taaagaacgt ggactccaac 4140
gtcaaagggc gaaaaaccgt ctatcagggc gatggcccac tacgtgaacc atcaccacaa 4200
tcaagttttt tgcggtcgag gtgccgtaaa gctctaaatc ggaaccctaa agggagcccc 4260
cgatttagag cttgacgggg aaagccggcg aacgtggcga gaaaggaagg gaagaaagcg 4320
aaaggagcgg gcgctagggc gctggcaagt gtagcggtca cgctgcgcgt aaccaccaca 4380
ccgcgcgcgc ttaatgcgcc gctacagggc gcgtccattc gccattcagg ctgcgcaact 4440
gttgggaagg gcgacgggtg cgggcctctt cgctattacg ccagctggcg aaagggggat 4500
gtgctgcaag gcgattaagt tgggtaacgc cagggttttc ccagtcacga cgttgtaaaa 4560
cgacggccag tgaattgtaa tacgactcac tataggcgga attcgagctc ggtacccggg 4620

```

<210> 3

<211> 4756

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Plasmid pGX22

<400> 3

| | | | | | | |
|-------------|-------------|------------|-------------|-------------|-------------|------|
| tgctcagaga | gtttctcaac | gaacccgatt | tggttagtta | taggtaattt | ttagaacatt | 60 |
| tacaaaaaca | gcaaaaaaac | caaacattca | ggatttttgt | ttttaattaa | gaaaaaaatc | 120 |
| gatcgctctt | aaattttaat | caatacttcg | aataaaacca | aaaaaaaacg | aaaaaaatc | 180 |
| ctgtttccag | tgtaatgatg | attgacgagg | ctcacgaacg | tactctacac | acggatattc | 240 |
| tattcggttt | agtcaaagat | attgcaagat | tccgaaagga | tttgaagctt | ctcatctctt | 300 |
| ctgcaacact | tgacgctgaa | aagttctcca | gtttcttcga | cgacgctccg | attttccgaa | 360 |
| ttccgggacg | cagattcccg | gtggacattt | actatacaca | ggctcccga | gcggactacg | 420 |
| tcgacgcggc | tatcgtcaca | attatgcaga | ttcacttgac | ccagccactt | ccggcgata | 480 |
| ttttggtatt | tctgactggt | caggaagaaa | tcgaaactgt | acaggaagca | cttatggaac | 540 |
| ggtcgaaagc | actgggatcg | aagattaagg | agcttattcc | gctgccggtt | tatgcgaatt | 600 |
| tgcccagtga | tttgcaggcg | aagattttcg | agccaacgcc | gaaagatgcg | agaaaggtag | 660 |
| atttttctta | caaatttttt | ccaaaaaaa | atccgagaaa | aatctacaaa | atttcaggca | 720 |
| aaaactgttt | catttttatc | ctaactagtt | ttttagcaaa | cgtttagatt | taacaaaact | 780 |
| gaacaaattt | gaagttttcc | aatttaaaaa | ataaatgttt | cggaaagtgt | attgaaaaat | 840 |
| ctgaaattgc | tatcctctcg | tatctgcaaa | aaaaacactt | taaaaaatgc | tctgttcttt | 900 |
| gaaaatttct | aaactgaaaa | atttgaaatt | tctgaaaatt | gtgataattt | tataaaattt | 960 |
| tatagaaaat | gtaagcattc | cagaaaaata | tcaaaaattt | cgagaaaatt | ctgaaaaaat | 1020 |
| ccagaaatat | taacagaaaa | aaaatctttt | gaaacatctg | aaaattaaaa | taaattgaat | 1080 |
| ttacattttt | ttttttggga | tttccttaaa | atcactatga | atttaccact | aaattttttg | 1140 |
| caaaaaatta | tttttttaat | ttcaaaagaa | aagcaaaagaa | ttttaaaata | tcaaaaagtc | 1200 |
| caaatttggt | tcggtgaaat | tttaaaataa | cattttcaag | ataattttta | gttaatcaaa | 1260 |
| acattccacg | catttctagt | ttcccaaatt | tctctaaatt | tcaggtggtc | ctagcaacta | 1320 |
| acattgccag | cacaatggat | ctcgagggat | cttccatacc | taccagttct | gcgcctgcag | 1380 |
| gtcgcggccg | cgactctcta | gacgcgtaag | cttactagca | taaccctctg | gggcctctaa | 1440 |
| acgggtcttg | aggggttttt | tgagcttctc | gccctatagt | gagtcgtatt | acagcttgag | 1500 |
| tattctatag | tgtcacctaa | atagcttggc | gtaatcatgg | tcatactgtg | ttcctgtgtg | 1560 |
| aaattgttat | cgcgtcacaa | ttccacacaa | catacgagcc | ggaagcataa | agtgtaaagc | 1620 |
| ctgggggtgcc | taatgagtga | gctaactcac | attaattgcg | ttgcgctcac | tgcccgtttt | 1680 |
| ccagtcggga | aaactgtcgt | gccagctgca | ttaatgaatc | ggccaacgcg | cggggagagg | 1740 |
| cggtttgcgt | attggggcgt | cttccgcttc | ctcgtcact | gactcgtgc | gctcggctcg | 1800 |
| tcggctgcgg | cgagcgggat | cagctcactc | aaaggcggta | atacggttat | ccacagaatc | 1860 |
| aggggataac | gcaggaaaaga | acatgtgagc | aaaaggccag | caaaaggcca | ggaaccgtaa | 1920 |
| aaaggccgcg | ttgctggcgt | ttttcgatag | gctccgcccc | cctgacgagc | atcacaaaaa | 1980 |
| tcgacgctca | agtcagaggt | ggcgaaaccc | gacaggacta | taaagatacc | aggcgtttcc | 2040 |
| ccctggaagc | tcctcgtgtc | gctctcctgt | tccgaccctg | ccgcttaccg | gatacctgtc | 2100 |
| cgcctttctc | ccttcgggaa | gcgtggcgct | ttctcatagc | tcacgctgta | ggtatctcag | 2160 |
| ttcgtgtgat | gtcgttcgct | ccaagctggg | ctgtgtgcac | gaaccccccg | ttcagcccca | 2220 |
| ccgctgcgcc | ttatccggta | actatcgtct | tgagccaac | ccggtaagac | acgacttata | 2280 |
| gccactggca | gcagccactg | gtaacaggat | tagcagagcg | aggatgtag | gcggtgctac | 2340 |
| agagttcttg | aagtgggtggc | ctaactacgg | ctacactaga | aggacagtat | ttggtatctg | 2400 |
| cgctctgctg | aagccagtta | ccttcggaaa | aagagttggt | agctcttgat | ccggcaaaaca | 2460 |
| aaccaccgct | ggtagcggtg | gtttttttgt | ttgcaagcag | cagattacgc | gcagaaaaaa | 2520 |
| aggatctcaa | gaagatcctt | tgatcttttc | tacggggtct | gacgctcagt | ggaacgaaaa | 2580 |
| ctcacgttaa | gggatttttg | tcatgagatt | atcaaaaagg | atcttcacct | agatcctttt | 2640 |
| aaattaaaaa | tgaagtttta | aatcaatcta | aagtatata | gagtaaaact | ggtctgacag | 2700 |
| ttaccaatgc | ttaatcagtg | aggcacctat | ctcagcgatc | tgtctatttc | gttcatccat | 2760 |
| agttgcctga | ctccccgtcg | tgtagataac | tacgatacgg | gagggcttac | catctggccc | 2820 |
| cagtgtctga | atgataccgc | gagaccacg | ctcaccggct | ccagatttat | cagcaataaa | 2880 |
| ccagccagcc | ggaaggggcg | agcgagaag | tggtcctgca | actttatccg | cctccatcca | 2940 |
| gtctattaat | gtttgcccgg | aagctagagt | aagtagttcg | ccagtttaata | gtttgcgcaa | 3000 |
| cgttgtttgg | attgctacag | gcacgtgggt | gtcacgctcg | tcgtttggta | tggttctcatt | 3060 |
| cagctccggt | tccaacgat | caaggcgagt | tacatgatcc | cccatgttgt | gcaaaaaagc | 3120 |
| ggttagctcc | ttcggctctc | cgatcgttgt | cagaagtaag | ttggccgcag | tgttatcact | 3180 |
| catggttatg | gcagcactgc | ataattctct | tactgtcatg | ccatccgtaa | gatgcttttc | 3240 |
| tgtgactggt | gagtactcaa | ccaagtcatt | ctgagaatac | cgcgccgggc | gaccgagttg | 3300 |
| ctcttgcccg | gcgtcaatac | gggataatag | tgtatgacat | agcagaactt | taaaagtgtc | 3360 |

```

catcattgga aaacgtttott cggggcgaaa actctcaagg atcttaccgc tgttgagatc 3420
cagttcgaag taacccactc gtgcacccaa ctgatcttca gcatctttta ctttcaccag 3480
cgtttctggg tgagcaaaaa caggaaaggca aaatgccgca aaaaaggga taaggcgac 3540
acggaaatgt tgaatactca tactcttcct ttttcaatat tattgaagca tttatcaggg 3600
ttattgtctc atgagcggat acatatattga atgtatttag aaaaataaac aaataggggt 3660
tccgcgcaca ttcccccga aagtgccacc tgacgtctaa gaaaccatta ttatcatgac 3720
attaacctat aaaaataggc gtatcacgag gccctttcgt ctgcgcggtt tcggtgatga 3780
cgggtgaaaac ctctgacaca tgcagctccc ggagacggtc acagcttgct tgtaagcgga 3840
tgccgggagc agacaagccc gtcaggggcg gtcagcgggt gttggcgggt gtcggggctg 3900
gcttaactat gcggcatcag agcagattgt actgagagt caccatatgc ggtgtgaaat 3960
accgcacaga tgcgtaagga gaaaataccg catcaggcga aattgtaaac gttaatattt 4020
tggtaaaatt cgcgttaaat atttgttaaa tcagctcatt ttttaacaa taggccgaaa 4080
tcggcaaaat cccttataaa tcaaaagaat agaccgagat agggttgagt gttgttccag 4140
tttggaacaa gagtccacta ttaaagaacg tggactcaa cgtcaaagg cgaaaaaccg 4200
tctatcaggc cgatggccca ctacgtgaac catcacccaa atcaagtttt ttgcggctga 4260
ggtgccgtaa agctctaaat cggaacccta aaggagccc ccgatttaga gcttgacggg 4320
gaaagccggc gaacgtggcg agaaaggagc ggaagaaagc gaaaggagcg ggcgctaggg 4380
cgctggcaag tctagcggtc acgctgcgcg taaccaccac acccgccgcg cttaatgcgc 4440
cgctacaggc cgcgtccatt cgccattcag gctgcgcaac tgttggaag ggcgatcggg 4500
gcgggcctct tcgctattac gccagctggc gaaaggggga tgtgctgcaa ggcgattaag 4560
ttgggtaaog ccagggtttt cccagtcacg acgttgtaaa acgacggcca gtgaattgta 4620
atacgactca ctatagggcg aattcaaaaa acccctcaag acccgtttag agggcccaag 4680
gggttatgct agtgaattct gcagggtacc cggggatcct ctagagatcc ctcgacctcg 4740
agatccattg tgctgg

```

<210> 4

<211> 4643

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Plasmid pGX52

<400> 4

```

gagtgcacca tatgcgggtgt gaaataccgc acagatgcgt aaggagaaaa taccgcatca 60
ggcgaaattg taaacgttaa tattttgtta aaattcgcgt taaatatttg ttaaatcagc 120
tcatttttta accaataggc cgaaatcggc aaaatccctt ataatcaaa agaataagacc 180
gagatagggt tgagtgttgt tccagtttg aacaagagtc cactattaaa gaacgtggac 240
tccaacgtca aagggcgaaa aaccgtctat caggggcgat gccactacg tgaacctatca 300
cccaaatcaa gttttttgcg gtcgaggtgc cgtaaagctc taaatcgaa ccctaaaggg 360
agccccgat ttagagcttg acggggaaag ccggcgaaag tggcgagaaa ggaagggaag 420
aaagcgaaag gagcgggcgc tagggcgctg gcaagtgtag cggtcacgct gcgcgtaacc 480
accacacccg ccgcgcttaa tgcgcgcta cagggcgcgt ccattcgcca ttcaggctgc 540
gcaactgttg ggaaggcgca tcggtgcggg cctcttcgct attacgccag ctggcgaaaag 600
ggggatgtgc tgcaaggcga ttaagttggg taacgccagg gttttccag tcacgacgtt 660
gtaaaacgac gccagtgaa ttgtaatac actcactata gggcgaattc aaaaaacccc 720
tcaagacccg ttagaggcc ccaaggggtt atgctagtga attctgcagg gtacccgggg 780
atcctctaga gatccctcga cctcgagatc cattgtgctg gcagccgatc tccgtcttgt 840
gaagatctac tccaccacca tccgtatcga tcagtccatc ctacccggag aatctgtgtc 900
tggtatcaag cacaccgact ctgtgccaga tccacgcgct gtaaccagg acaagaagaa 960
ttgtctgttc tcgggaacca atgtcgcac tggaaaggct cgtggaatcg tcttcggaac 1020
cggattgacc actgaaatcg gaaagatccg taccgaaatg gctgagaccg agaatgagaa 1080
gacaccactt caacagaagt tggacgaatt cggagagcaa ctttccaagg ttatctctgt 1140
tatttgctgt gctgtttggg ctatcaacat tggacatttc aacgatccag ctacggtggg 1200
atcatgggtt aaggagcaa tctactactt caaaatcgcc gttgctcttg ccgtcgctgc 1260
tattccagaa ggaactccag ctgtcatcac cacgtgcctt gccctcgaa ctgcgctgat 1320
ggccaagaag aacgctattg taagatccct tccatccgtc gaaactcttg gatgcacatc 1380
tggtatctgc tctgacaaga ctggaactct caccaccaac cagatgtctg tgtcaaagat 1440

```

| | | | | | | |
|-------------|------------|-------------|------------|-------------|-------------|------|
| gttcacgcgt | ggacaagctt | ctggagacaa | catcaacttc | accgagttcg | ccatctccgg | 1500 |
| atccacctac | gagccagtcg | gaaaggtttc | caccaatgga | cgtgaaatca | acccagctgc | 1560 |
| tggagaattc | gaatcactca | ccgagttggc | catgatctgc | gctatgtgca | atgattcatc | 1620 |
| tgttgattac | aatgagacca | agaagatcta | cgagaaagtc | ggagaagcca | ctgaaactgc | 1680 |
| tcttatcggt | cttgctgaga | agatgaatgt | tttcggaacc | tcgaaagccg | gactttcacc | 1740 |
| aaaggagctc | ggaggagttt | gcaaccgtgt | catccaaaca | aaatggaaga | aggagttcac | 1800 |
| actcgagttc | tcccgtgata | gtaaataccat | gtccgcctac | tgcttccag | cttccggagg | 1860 |
| atctggagcc | aagatgttcg | tgaaggagc | cccagaagga | gttctcggaa | gatgcaccca | 1920 |
| cgtcagagtt | aacggacaaa | aggttccact | cacctctgcc | atgactcaga | agattgttga | 1980 |
| ccaatgcgtg | caatacggaa | ccggaagaga | tacccttcgt | tgtcttgccc | tcggccagca | 2040 |
| caatggatct | cgaggatctc | tccataccta | ccagttctgc | gcctgcaggt | cgcgcccgcg | 2100 |
| actctctaga | cgcgtaagct | tactagcata | accccttggg | gcctctaaac | gggtcttgag | 2160 |
| gggttttttg | agcttctcgc | cctatagtga | gtcgtattac | agcttgagta | ttctatagtg | 2220 |
| tcacctaaat | agcttggcgt | aatcatggtc | atagctgttt | cctgtgtgaa | attgttatcc | 2280 |
| gctcacaatt | ccacacaaca | tacgagccgg | aagcataaag | tgtaaagcct | ggggtgccta | 2340 |
| atgagtgaac | taactcacat | taattgcgtt | gcgctcactg | cccgttttcc | agtcgggaaa | 2400 |
| cctgtcgtgc | cagctgcatt | aatgaatcgg | ccaacgcgcg | gggagagggc | gtttgcgtat | 2460 |
| tggcgctctc | tccgcttcc | cgctcactga | ctcgtcgcgc | tcggtcgttc | ggctgcggcg | 2520 |
| agcggatatca | gctcactcaa | aggcggtaat | acggttatcc | acagaatcag | gggataacgc | 2580 |
| aggaaagaac | atgtgagcaa | aaggccagca | aaaggccagg | aaccgtaaaa | aggccgcggt | 2640 |
| gctggcggtt | ttcgatagga | tccgcccccc | tgacgagcat | cacaaaaatc | gacgctcaag | 2700 |
| tcagaggtgg | cgaaacccga | caggactata | aagataccag | gcgtttcccc | ctggaagctc | 2760 |
| cctcgtgcgc | tctcctgttc | cgaccctgcc | gcttaccgga | tacctgtccg | cctttctccc | 2820 |
| ttcgggaagc | gtggcgcttt | ctcatagctc | acgctgtagg | tatctcagtt | cggtgtagg | 2880 |
| cgttcgctcc | aagctgggct | gtgtgcacga | acccccggtt | cagcccgacc | gctgcgcctt | 2940 |
| atccggtaac | tatcgtcttg | agtccaaccc | ggtaagacac | gacttatcgc | cactggcagc | 3000 |
| agccactggt | aacaggatta | gcagagcgag | gtatgtagcc | gggtgtacag | agttcttgaa | 3060 |
| gtggtggcct | aactacggct | acactagaag | gacagtattt | ggatatctgc | ctctgctgaa | 3120 |
| gccacttacc | ttcggaaaaa | gagttggtag | ctcttgatcc | ggcaaacaaa | ccaccgctgg | 3180 |
| tagcgggtgt | ttttttgttt | gcaagcagca | gattacgcgc | agaaaaaaag | gatctcaaga | 3240 |
| agatcctttg | atcttttcta | cggggtctga | cgctcagtg | aacgaaaact | cacgttaagg | 3300 |
| gattttggtc | atgagattat | caaaaaggat | cttcacctag | atccttttaa | attaaaaatg | 3360 |
| aagttttaaa | tcaatctaaa | gtatatatga | gtaaacttgg | tctgacagtt | accaatgctt | 3420 |
| aatcagtgag | gcacctatct | cagcgatctg | tctatttcgt | tcattccatag | ttgcctgact | 3480 |
| ccccgtcgtg | tagataacta | cgatacggga | gggcttacca | tctggcccca | gtgctgcaat | 3540 |
| gataccgcga | gaccacgcgt | caccggctcc | agatttatca | gcaataaacc | agccagccgg | 3600 |
| aagggccgag | cgcagaagtg | gtcctgcaac | tttatccgcc | tccatccagt | ctattaattg | 3660 |
| ttgcccggaa | gctagagtaa | gtagtccgcc | agttaatagt | ttgcgcaacg | ttgttggtat | 3720 |
| tgctacagga | atcgtgggtg | cacgctcgtc | gtttggtag | gcttcattca | gctccgggtc | 3780 |
| ccaacgatca | aggcgagtta | catgatcccc | catgttggtc | aaaaaagcgg | ttagctcctt | 3840 |
| cggctctacc | atcgttggtc | gaagtaagtt | ggcgcgagtg | ttatcactca | tgggttatggc | 3900 |
| agcactgcat | aattctctta | ctgtcatgcc | atccgtaaga | tgcttttctg | tgactggtga | 3960 |
| gtactcaacc | aagtcattct | gagaataccg | cgcccgccga | ccgagttgct | cttgcccggc | 4020 |
| gtcaatacgg | gataatagt | tatgacatag | cagaacttta | aaagtgtca | tcattggaaa | 4080 |
| acgttcttcg | ggcgaaaaac | tctcaaggat | cttaccgctg | ttgagatcca | gttcgatgta | 4140 |
| acccactcgt | gcacccaact | gatcttcagc | atcttttact | ttcaccagcg | tttctgggtg | 4200 |
| agcaaaaaa | ggaaggcaaa | atgccgcaaa | aaagggaata | agggcgacac | ggaaatggtg | 4260 |
| aatactcata | ctcttccctt | ttcaatatta | ttgaagcatt | tatcagggtt | attgtctcat | 4320 |
| gagcggatac | atatttgaat | gtatttagaa | aaataaaca | ataggggttc | cgcgcacatt | 4380 |
| tccccgaaaa | gtgccacctg | acgtctaaga | aaccattatt | atcatgacat | taacctataa | 4440 |
| aaataggcgt | atcacgaggc | cctttcgtct | cgcgcgtttc | ggtgatgacg | gtgaaaacct | 4500 |
| ctgacacatg | cagctcccgg | agacggteac | agcttctctg | taagcggatg | ccgggagcag | 4560 |
| acaagccgct | caggggcgct | cagcgggtgt | tggcgggtgt | cggggctggc | ttactatgct | 4620 |
| ggcatcagag | cagattgtac | tga | | | | 4643 |

<210> 5

<211> 4454

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Plasmid pGX104

<400> 5

| | | | | | | |
|-------------|-------------|-------------|------------|------------|-------------|------|
| gagtgcacca | tatgcggtgt | gaaataccgc | acagatgcgt | aaggagaaaa | taccgcatca | 60 |
| ggcgaaattg | taaacgttaa | tattttgtta | aaattcgcgt | taaatatttg | ttaaatacagc | 120 |
| tcatttttta | accaataggc | cgaaatcggc | aaaatcccct | ataaatcaaa | agaatagacc | 180 |
| gagataggg | tgagtgttgt | tccagtttgg | aacaagagtc | cactatttaa | gaacgtggac | 240 |
| tccaacgtca | aagggcgaaa | aaccgtctat | cagggcgatg | gcccactacg | tgaaccatca | 300 |
| cccaaatcaa | gttttttgcg | gtcgaggtgc | cgtaaagctc | taaatcgga | ccctaaaggg | 360 |
| agccccgat | ttagagcttg | acggggaaag | ccggcgaaag | tggcgagaaa | ggaagggag | 420 |
| aaagcgaaag | gagcggggcg | tagggcgctg | gcaagtgtag | cggtcacgct | gcgcgtaacc | 480 |
| accacacccg | ccgcgcttaa | tgcccgctta | cagggcgctg | ccattcgcca | ttcaggtctg | 540 |
| gcaactgttg | ggaagggcga | tcggtgcggg | cctcttcgct | attacgccag | ctggcgaaag | 600 |
| ggggatgtgc | tgcaaggcga | ttaagtgtgg | taacgccagg | gttttcccag | tcacgacgtt | 660 |
| gtaaaacgac | ggccagtga | ttgtaatacg | actcactata | gggcgaattc | aaaaaacccc | 720 |
| tcaagaccgg | tttagaggcc | ccaaggggtt | atgctagtga | attctgcagg | gtacccgggg | 780 |
| atcctctaga | gatccctcga | cctcgagatc | cattgtgctg | gaccgtggta | ctcttatgga | 840 |
| gctcggaatc | tcgccaatcg | tcacttctgg | acttatcatg | caacttctcg | ccggagccaa | 900 |
| gatcatcgaa | gtcggagaca | caccaaagga | ccgtgctctt | ttcaacggag | cccagaaatg | 960 |
| taagccgaaa | agtgtgtgtt | ttcaatctct | aatttttgaa | cttttcagtg | ttcgggtatg | 1020 |
| tcactactgt | tggacaagct | attgtctacg | tcattgtccg | actctacgga | gagccatcgg | 1080 |
| aaatcgagc | tggaaatctg | ctccttatcg | tcgtccaact | cgttattgcc | ggtctcatcg | 1140 |
| tcctccttct | cgacgagctt | ctccaaaagg | gatatggtct | cggtatccga | atttctctct | 1200 |
| tcattgccac | caacatctgt | gaaaccattg | tctggaaggc | attctccccg | gcaacaatga | 1260 |
| acacgggacg | tggaaaccgag | ttcgaaggag | ccgtcattgc | tcttttccat | cttcttgcca | 1320 |
| cccgtccga | caaggtccgt | gcccttcgtg | aggctttcta | ccgtcaaaac | cttccaaact | 1380 |
| tgatgaactt | gatggctact | ttcctcggtt | ttgcggtggt | tatctacttc | caaggattcc | 1440 |
| gtgtcgacct | cccaatcaag | tctgcccgct | accgtggaca | atacagcagc | taccaatca | 1500 |
| agctcttcta | cacctccaac | attccaatca | tccttcaatc | tgctctcgct | tccaacctct | 1560 |
| acgttatctc | tcaggtttgt | tgcatctcag | tagtacggtt | agatgtttat | ctttctctag | 1620 |
| agggctcaagt | tggcggagaa | attttttgag | ttcattctca | agtctgatgg | aaaatgttta | 1680 |
| tttttcagat | gtcgcggga | aagttcggag | gaaacttctt | catcaacctt | ctcgggtacct | 1740 |
| ggtccgataa | caccggatac | agaagctacc | caactggagg | actctgctac | tatctttcac | 1800 |
| caccagagtc | tcttggaac | atcttcgaag | acccaatcca | ctgcaccagc | acaatggatc | 1860 |
| tcgagggatc | ttccatacct | accagtctct | cgcttcgagg | tcgcggcgcg | gactctctag | 1920 |
| acgcgtatgc | tactagcat | aacccttggt | ggcctctaaa | cgggtcttga | ggggtttttt | 1980 |
| gagcttctcg | ccctatagtg | agtcgtatta | cagcttgagt | attctatagt | gtcacctaaa | 2040 |
| tagcttggcg | taatcatggt | catagctggt | tcctgtgtga | aattgtttat | cgctcacaat | 2100 |
| tccacacaac | atacagaccg | gaagcataaa | gtgtaaagcc | tgggggtgct | aatgagtga | 2160 |
| ctaactcaca | ttaattgctg | tgcgctcact | gcccgctttc | cagtcgggaa | acctgtcggt | 2220 |
| ccagctgcat | taatgaatcg | gccaacgcgc | ggggagaggg | ggtttgcgta | ttggggcgct | 2280 |
| ttccgcttcc | tcgctcactg | actcgctgcg | ctcggtcggt | cggctgcggc | gagcgggtat | 2340 |
| agctcactca | aaggcggtaa | tacggttatc | cacagaatca | ggggataacg | caggaaagaa | 2400 |
| catgtgagca | aaaggccagc | aaaaggccag | gaaccgtaaa | aaggccgcgt | tgctggcggt | 2460 |
| tttcgatagg | ctccgcccc | ctgacgagca | tcacaaaaat | cgacgctcaa | gtcagagggt | 2520 |
| gcgaaacccg | acaggactat | aaagatacca | ggcggtttcc | cctggaagct | ccctcggtcg | 2580 |
| ctctcctggt | ccgaccctgc | cgcttaccgg | atacctgtcc | gcctttctcc | cttcgggaag | 2640 |
| cgtggcgctt | tctcatagct | cagctgttag | gtatctcagt | tcggtgtagg | tcgttcgctc | 2700 |
| caagctgggc | tgtgtgcacg | aacccccgt | tcagcccgc | cgctgcgcct | tatccggtaa | 2760 |
| ctatcgtctt | gagtccaacc | cggtaaagaca | cgacttatcg | ccactggcag | cagccactgg | 2820 |
| taacaggatt | agcagagcga | ggtatgtagg | cggtgctaca | gagttcttga | agtgggtggc | 2880 |
| taactacggc | tacactagaa | ggacagtatt | tggtatctgc | gctctgctga | agccagttac | 2940 |
| cttcggaaaa | agagttggta | gctcttgatc | cggcaaaaa | accaccgctg | gtagcgggtg | 3000 |
| tttttttgtt | tgcaagcagc | agattacgcg | cagaaaaaaa | ggatctcaag | aagatccttt | 3060 |
| gatcttttct | acgggggtctg | acgctcagtg | gaacgaaaa | tcacgttaag | ggattttggt | 3120 |

8

```

catgagatta tcaaaaagga tcttcaccta gatcctttta aattaaaaat gaagttttta 3180
atcaatctaa agtatatatg agtaaaacttg gtctgacagt taccaatgct taatcagtga 3240
ggcacctatc tcagcgatct gtctatttcg ttcattccata gttgcctgac tccccgctgt 3300
gtagataact acgatacggg agggccttacc atctggcccc agtgctgcaa tgataccgcg 3360
agaccacgc tcaccggctc cagatttatc agcaataaac cagccagccg gaagggccga 3420
gcgcagaagt ggtcctgcaa ctttatccgc ctccatccag tctattaatt gttgccggga 3480
agctagagta agtagttcgc cagttaatag tttgcgcaac gttgttgga ttgctacagg 3540
catcgtgggtg tcacgctcgt cgtttggtat ggcttcattc agctccggtt cccaacgatc 3600
aaggcgagtt acatgatccc ccatgttggtg caaaaaagcg gttagctcct tcgggtcctcc 3660
gatcgttggtc agaagtaagt tggccgcagt gttatcactc atgggttatgg cagcactgca 3720
taattctctt actgtcatgc catccgtaag atgcttttct gtgactgggt agtactcaac 3780
caagtcatte tgagaatacc gcgcccgcg accgagttgc tcttgccggg cgtcaatacg 3840
ggataaatagt gtatgacata gcagaacttt aaaagtgtc atcattggaa aacgttcttc 3900
ggggcgaaaa ctctcaagga tcttaccgct gttgagatcc agttcgatgt aaccactcg 3960
tgcacccaac tgatcttcag catcttttac tttcaccagc gtttctgggt gagcaaaaac 4020
aggaaggcaa aatgccgcaa aaaagggaa aagggcgaca cggaatggt gaatactcat 4080
actcttctct tttcaatatt attgaagcat ttatcagggt tattgtctca tgagcgata 4140
catatttgaa tgtatttaga aaaataaaca aatagggtt ccgcgcacat tccccgaaa 4200
agtgccacct gacgtctaag aaaccattat tatcatgaca ttaacctata aaaataggcg 4260
tatcacgagg ccctttcgtc tcgcgcgttt cgggtgatgc ggtgaaaacc tctgacacat 4320
gcagctcccg gagacgggtc cagcttgtct gtaagcggat gccgggagca gacaagccc 4380
tcagggcgcg tcagcgggtg ttggcgggtg tcggggctgg cttactatg cggcatcaga 4440
gcagattgta ctga                                     4454

```

<210> 6

<211> 4701

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Plasmid pGZ18

<400> 6

```

accagcttt cttgtacaaa gtggtgatct ttccagcaca atggatctcg aggatcttc 60
catactacc agttctgctc ctgcaggtcg oggcccgcac tctctagacg cgttaagctta 120
ctagcataac cccttggggc ctctaaacgg gtcttgaggg gttttttgag cttctcgccc 180
tatagtgagt cgtattacag cttgagtatt ctatagtgtc acctaaatag cttggcgtaa 240
tcatggtcat agctgtttcc tgtgtgaaat tgttatccgc tcacaattcc acacaacata 300
cgagccggaa gcataaagtg taaagcctgg ggtgccta at gagtgagcta actcacatta 360
attgcttgc gctcactgcc cgctttccag tcgggaaacc tgcgtgcca gctgcattaa 420
tgaatcgcc aacgcgcggg gagaggcgtt ttgcgtattg ggcgctcttc cgcttccctg 480
ctcactgact cgctgcgtc ggtcgttcgg ctgcggcgag cggtatcagc tcaactcaaag 540
gcggtaatat ggttatccac agaatcaggg gataacgcag gaaagaacat gtgagcaaaa 600
ggccagcaaa aggccaggaa ccgtaaaaag gccgcgttgc tggcgttttt cgataggctc 660
cgccccctg acgagcatca caaaaaatcga cgctcaagtc agaggtggcg aaaccgcaca 720
ggactataaa gataccagge gtttccccct ggaagctccc tcgtgcgtc tcctgttccg 780
accctgcgcg ttaccggata cctgtccgcc tttctccctt cgggaagcgt ggcgctttct 840
catagctcac gctgtaggta tctcagttcg gtgtaggtcg ttcgctccaa gctgggctgt 900
gtgcacgaac ccccgttca gcccgaccgc tgcgccttat ccggttaacta tcgtcttgag 960
tccaaccgg taagacacga cttatcgcca ctggcagcag ccaactggtaa caggattagc 1020
agagcgaggt atgtaggcgg tgctacagag ttcttgaagt ggtggcctaa ctacggctac 1080
actagaagga cagtatttgg tatctgcgt ctgctgaagc cagttacctt cggaataaga 1140
gttggtagct cttgatccgg caaacaacc accgctggta gcggtggttt ttttgtttgc 1200
aagcagcaga ttacgcgcag aaaaaagga tctcaagaag atcctttgat cttttctacg 1260
gggtctgacg ctcagtgga cgaaaactca cgtaaggga ttttgggtcat gagattatca 1320
aaaaggatct tcacctagat ctttttaaat taaaaatgaa gtttttaaat aatctaaagt 1380
atatatgagt aaacttggtc tgacagttac caatgcttaa tcagtgggc acctatctca 1440
gcgatctgtc tatttcgttc atccatagtt gcctgactcc ccgtcgtgta gataactacg 1500

```

| | | | | | | |
|-------------|-------------|------------|-------------|-------------|-------------|------|
| atacgggagg | gcttaccatc | tggccccagt | gctgcaatga | taccgcgaga | cccacgctca | 1560 |
| ccgggtccag | atztatcagc | aataaaccag | ccagccggaa | gggcccagcg | cagaagtggg | 1620 |
| cctgcaactt | tatccgcctc | catccagttc | attaattgtt | gccgggaagc | tagagtaagt | 1680 |
| agttcgcgag | ttaatatgtt | gcgcaacggt | gttggcattg | ctacaggcat | cgtgggtgtca | 1740 |
| cgctcgtcgt | ttggtatggc | ttcattcagc | tccggttccc | aacgatcaag | gcgagttaca | 1800 |
| tgatccccc | tggtgtgcaa | aaaagcgggt | agctccttcg | gtcctccgat | cggtgtcaga | 1860 |
| agtaagtgg | ccgcagtggt | atcactcatg | gttatggcag | cactgcataa | ttctcttact | 1920 |
| gtcatgccat | ccgtaagatg | cttttctgtg | actggtgagt | actcaaccaa | gtcattctga | 1980 |
| gaataccgcg | cccggcgacc | gagttgctct | tgccccggcg | caatacggga | taatagtgtg | 2040 |
| tgacatagca | gaacttttaa | agtgtctatc | attggaaaac | gttcttcggg | gcgaaaactc | 2100 |
| tcaaggatct | taccgctgtt | gagatccagt | tcgatgtaac | ccactcgtgc | acccaactga | 2160 |
| tcttcagcat | cttttacttt | caccagcggt | tctgggtgag | caaaaacagg | aaggcaaaat | 2220 |
| gccgcaaaaa | aggggaataag | ggcgacacgg | aaatgttgaa | tactcatact | cttccttttt | 2280 |
| caatattatt | gaagcattta | tcaggggtat | tgtctcatga | gcggatacat | atttgaatgt | 2340 |
| atttagaaaa | ataaacaat | aggggttccg | cgcacatttc | cccgaaaagt | gccacctgac | 2400 |
| gtctaagaaa | ccattattat | catgacatta | acctataaaa | ataggcggtat | cacgaggccc | 2460 |
| tttctgctcg | cgcgttttcg | tgatgacggt | gaaaacctct | gacacatgca | gctcccggag | 2520 |
| acggtcacag | cttggtctgta | agcggatgcc | gggagcagac | aagcccgta | ggcgcggtca | 2580 |
| gcgggtgttg | gcgggtgtcg | gggctggctt | aactatgcgg | catcagagca | gattgtactg | 2640 |
| agagtgcacc | atatgcggtg | tgaaataccg | cacagatgcg | taaggagaaa | ataccgcctc | 2700 |
| aggcgaaatt | gtaaacgtta | atattttgtt | aaaattcgcg | ttaaatattt | gttaaatcag | 2760 |
| ctcatttttt | aaccaatagg | ccgaaatcgg | caaaatccct | tataaatcaa | aagaatagac | 2820 |
| cgagatagg | ttgagtgttg | ttccagtttg | gaacaagagt | ccactattaa | agaacgtgga | 2880 |
| ctccaacgtc | aaagggcgaa | aaaccgtcta | tcagggcgat | ggcccactac | gtgaaccatc | 2940 |
| acccaaatca | agttttttgc | ggtcgaggtg | ccgtaaagct | ctaaatcgga | accctaaagg | 3000 |
| gagccccgga | tttagagctt | gacggggaaa | gccggcgaa | gtggcgagaa | aggaagggaa | 3060 |
| gaaagcgaaa | ggagcggg | ctagggcgct | ggcaagtgtg | gcggtcacgc | tgcgcgtaac | 3120 |
| caccacacc | gccgcgctta | atgcgcgct | acagggcgcg | tccattcgcc | atcaggctg | 3180 |
| cgcaactgtt | gggaagggcg | atcgggtcgg | gcctcttcgc | tattacgcca | gctggcgaaa | 3240 |
| gggggatgtg | ctgcaaggcg | attaagttgg | gtaacgccag | ggttttccca | gtcacgacgt | 3300 |
| tgtaaaacga | cggccagtg | attgtaatac | gactcactat | agggcggaatt | caaaaaaccc | 3360 |
| ctcaagaccc | gttttagagg | cccaaggggt | tatgctagt | aattctgcag | ggtacccggg | 3420 |
| gaccccttag | agatccctcg | acctcgagat | ccattgtgct | ggaaagcctt | tgcaaggctg | 3480 |
| gcaagccacg | tttgggtgtg | gcgaccatcc | tccaaaatca | acaagtttgt | acaaaaaagc | 3540 |
| aggctatgcc | aagtacatgt | cgattgcgta | cgcggttcgta | atgttggctg | tgtagtgcgc | 3600 |
| taccagcagt | caaattgttc | tcgagagtgc | gtttttacat | tatcccttca | tcctgattac | 3660 |
| gacaattttc | agctgtttct | gtcctacat | ctctcttcat | tgtcacaatg | gtcggaaatct | 3720 |
| tcttctttgc | tgcatgtctt | catccaaaag | aattcacgaa | tattatccat | ggtgtcgtat | 3780 |
| tcttctctcat | gattccatct | acatatgtgt | tcctcacttt | atattcgctc | atcaatctca | 3840 |
| acgttatcac | gtggggaaact | cgtgaagctg | tcgctaagcg | aacgggacaa | aagacgaaaa | 3900 |
| aagcgcctat | ggaacaattt | atagacagag | tgattgatat | tgtgaaaaag | ggattcagat | 3960 |
| taatcagttg | tcgggagaag | aaggacatg | aagagagacg | agagaaaatg | gaaaagaaaa | 4020 |
| tgagagaaat | ggagctagcc | ttgagaagta | ttgaggttat | ctttaacttt | agaaatgtga | 4080 |
| aattaataat | ttattttcag | agtgggtccg | acgtgaagaa | aattctcgat | gcaacagagg | 4140 |
| agaaggagaa | acgtgaagaa | gaaactcaaa | tcgcagattt | tccgattgaa | gagaacgtag | 4200 |
| agaagactca | aaaagagatt | cagaaggcaa | accgttatgt | gtggatgaca | agtcataagc | 4260 |
| tgaaagtgtg | tgaacgagga | aaactgaaaa | gtgcggaaaa | ggttttctgg | aacgagctca | 4320 |
| tcaatgcata | tctgaaaccg | atcaagacga | cgccagctga | aatgaaagcc | gtcgccgaag | 4380 |
| gattggcttc | tctacgaaat | cagattgctt | tcactattct | tctcgtaaat | tctcttcttg | 4440 |
| ctcttgccat | ctttttgatt | cagaaacaca | aaaatgtgct | cagcatcaag | ttctcgccaa | 4500 |
| tcagtaagca | atattacctt | tatggtaaat | tcaaaaaatt | tggttttttt | ttctagaaaa | 4560 |
| cttccgatgg | acgaaaatga | atgagatgac | tggaacaatac | gaggaaaaccg | atgaaccatt | 4620 |
| aaaaatagat | ccacttgga | tggaattgt | tggttttctt | ctaattattc | tttttgttca | 4680 |
| aactctcgga | atgcttctcc | a | | | | 4701 |

<210> 7

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
Oligonucleotide primer C04H5.6F

<400> 7

tgctcagaga gtttctcaac gaacc

25

<210> 8

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Oligonucleotide
primer C04H5.6R

<400> 8

caatgttagt tgctaggacc acctg

25

<210> 9

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Oligonucleotide
primer K11D9.2bF

<400> 9

cagccgatct cagtcttgtg

20

<210> 10

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Oligonucleotide
primer K11D9.2bR

<400> 10

ccgagggcaa gacaacgaag

20

<210> 11

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Oligonucleotide
primer Y57G11C.15F

<400> 11

accgtggtac tcttatggag ctcg

24

<210> 12

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Oligonucleotide
primer Y57G11C.15R

<400> 12

tgcaagtggat tgggtcttcg

20

<210> 13

<211> 52

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Oligonucleotide
primer T25G3.2F

<400> 13

ggggacaagt ttgtacaaaa aagcaggcta tgccaagtac atgtcgattg cg

52

<210> 14

<211> 52

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Oligonucleotide
primer T25G3.2R

<400> 14

ggggaccact ttgtacaaga aagctggggtt ggagaagcat tccgagagtt tg

52

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/10 C12N9/22 C07K14/435 C12N15/66 C12N15/70
C12N1/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

BIOSIS, EPO-Internal, WPI Data, PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category * | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|------------|---|---|
| X | TIMMONS L ET AL: "Specific interference by ingested dsRNA" NATURE, MACMILLAN JOURNALS LTD. LONDON, GB, vol. 395, no. 6705, 29 October 1998 (1998-10-29), page 854 XP002103601 ISSN: 0028-0836 the whole document | 1-4, 12-21, 23-25, 28-31, 34-39 |
| A | FIRE A ET AL: "Potent and specific genetic interference by double-stranded RNA in Caenorhabditis elegans" NATURE, GB, MACMILLAN JOURNALS LTD. LONDON, vol. 391, 19 February 1998 (1998-02-19), pages 806-811, XP002095876 ISSN: 0028-0836 cited in the application the whole document | 1-4 |

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

G document member of the same patent family

Date of the actual completion of the international search

5 July 2001

Date of mailing of the international search report

20/07/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Mateo Rosell, A.M.

| C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT | | |
|--|--|-----------------------|
| Category * | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
| A | SHARP PHILLIP A: "RNAi and double-strand RNA." GENES & DEVELOPMENT, vol. 13, no. 2, 15 January 1999 (1999-01-15), pages 139-141, XP002171268 ISSN: 0890-9369 the whole document --- | 1-4 |
| A | RAY C ET AL: "GUT-SPECIFIC AND DEVELOPMENTAL EXPRESSION OF A CAENORHABDITIS ELEGANS CYSTEINE PROTEASE GENE" MOLECULAR AND BIOCHEMICAL PARASITOLOGY,NL,ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM, vol. 51, 1992, pages 239-249, XP000572340 ISSN: 0166-6851 abstract --- | 2-4,8 |
| A | RAND J B ET AL: "GENETIC PHARMACOLOGY: INTERACTIONS BETWEEN DRUGS AND GENE PRODUCTS IN CAENORHABDITIS ELEGANS" METHODS IN CELL BIOLOGY,LONDON,GB, vol. 84, 1995, pages 187-204, XP000956211 page 190, paragraph 1 -page 194, paragraph 4 --- | 1-4,8 |
| A | AVERY LEON ET AL: "The Caenorhabditis elegans unc-31 gene affects multiple nervous system-controlled functions." GENETICS, vol. 134, no. 2, 1993, pages 455-464, XP001011453 ISSN: 0016-6731 the whole document --- | 2-4,8 |
| A | TAGESSON C ET AL: "INFLUENCE OF SURFACE-ACTIVE FOOD ADDITIVES ON THE INTEGRITY AND PERMEABILITY OF RAT INTESTINAL MUCOSA" FOOD AND CHEMICAL TOXICOLOGY, vol. 22, no. 11, 1984, pages 861-864, XP001009621 ISSN: 0278-6915 the whole document --- -/-- | 8 |

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

| Category * | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|------------|--|---|
| P,A | <p>LYON CHRISTOPHER J ET AL: "The C. elegans apoptotic nuclease NUC-1 is related in sequence and activity to mammalian DNase II."</p> <p>GENE (AMSTERDAM), vol. 252, no. 1-2, 11 July 2000 (2000-07-11), pages 147-154, XP001009494 ISSN: 0378-1119 abstract; figure 3 page 151, right-hand column, paragraph 2 -page 153, right-hand column, paragraph 2</p> <p>---</p> | 6,7 |
| P,X | <p>WO 00 01846 A (MORTIER KATHERINE ;DEVGEN NV (BE); BOGAERT THIERRY (BE); PLAETINCK) 13 January 2000 (2000-01-13) cited in the application</p> <p>page 8, line 9 -page 10, line 22 page 12, line 14 -page 23, line 2 figures 5,9; examples 1-4</p> <p>---</p> | 1-4, 12-21, 23-25, 28-31, 34-39 |
| P,X | <p>WO 00 63425 A (FEICHTINGER RICHARD ;BEGHYN MYRIAM (BE); DEVGEN NV (BE); BOGAERT T) 26 October 2000 (2000-10-26) abstract page 2, line 14-30 page 6, line 18-33 page 7, line 12 -page 9, line 28 page 12, line 20 -page 13, line 25; example 3</p> <p>---</p> | 2-5,8,9, 29-33 |
| T | <p>TIMMONS LISA ET AL: "Ingestion of bacterially expressed dsRNAs can produce specific and potent genetic interference in Caenorhabditis elegans."</p> <p>GENE (AMSTERDAM), vol. 263, no. 1-2, 2001, pages 103-112, XP001009512 ISSN: 0378-1119 the whole document</p> <p>-----</p> | 1-4, 12-21, 23-26, 28-31, 34-39 |

| Patent document cited in search report | Publication date | Patent family member(s) | Publication date |
|---|---------------------|----------------------------|---------------------|
| WO 0001846 A | 13-01-2000 | AU 4907999 A | 24-01-2000 |
| | | EP 1093526 A | 25-04-2001 |
| | | GB 2349885 A | 15-11-2000 |
| | | NO 20010019 A | 05-03-2001 |
| <hr/> | | | |
| WO 0063425 A | 26-10-2000 | AU 3984600 A | 02-11-2000 |
| | | GB 2351152 A | 20-12-2000 |
| <hr/> | | | |

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.